

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) **EP 0 874 045 A1**

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

28.10.1998 Bulletin 1998/44

(51) Int. Cl.⁶: **C12N 15/00**, C12P 21/00

(21) Application number: **97935810.8**

(86) International application number:

PCT/JP97/02859

(22) Date of filing: **19.08.1997**

(87) International publication number:

WO 98/07840 (26.02.1998 Gazette 1998/08)

(84) Designated Contracting States:

AT BE CH DE DK ES FR GB IE IT LI LU NL SE

(30) Priority: **19.08.1996 JP 235928/96**

(83) Declaration under Rule 28(4) EPC (expert solution)

(71) Applicant:

SNOW BRAND MILK PRODUCTS CO., LTD.
Sapporo-shi, Hokkaido 065 (JP)

(72) Inventors:

- **NAKAGAWA, Nobuaki,**
Nishiura Heights 2-4
Shimotsuga-gun, Tochigi 329-05 (JP)
- **YASUDA, Hisataka**
Kawachi-gun, Tochigi 329-04 (JP)
- **MORINAGA, Tomonori**
Shimotsuga-gun, Tochigi 321-02 (JP)

(74) Representative:

Wakerley, Helen Rachael
Reddie & Grose,
16 Theobalds Road
London WC1X 8PL (GB)

(54) **NOVEL DNAS AND PROCESS FOR PRODUCING PROTEINS BY USING THE SAME**

(57) DNAs having the nucleotide sequences of the Sequences No. 1 and No. 2 in the Sequence Table and a process for producing a protein which comprises inserting these DNAs into expression vectors to thereby produce a protein having molecular weights of about 60 kD (under reductive conditions) and about 60 kD and 120 kD (under non-reductive conditions) and being capable of inhibiting formation of osteoclast. These proteins are useful in the treatment of osteoporosis and rheumatism.

EP 0 874 045 A1

Description

FIELD OF TECHNOLOGY

5 The present invention relates to a novel DNA and a process for preparing a protein which possesses an activity to inhibit osteoclast differentiation and/or maturation (hereinafter called osteoclastogenesis-inhibitory activity) by a genetic engineering technique using the DNA. More particularly, the present invention relates to a genomic DNA encoding a protein OCIF which possesses an osteoclastogenesis-inhibitory activity and a process for preparing said protein by a genetic engineering technique using the genomic DNA.

BACKGROUND OF THE INVENTION

10 Human bones are constantly repeating a process of resorption and formation. Osteoblasts controlling formation of bones and osteoclasts controlling resorption of bones take major roles in this process. Osteoporosis is a typical disease caused by abnormal metabolism of bones. This disease is caused when bone resorption by osteoclasts exceeds bone formation by osteoblasts. Although the mechanism of this disease is still to be elucidated completely, the disease causes the bones to ache, makes the bones fragile, and may results in fracturing of the bones. As the population of the aged increases, this disease results in an increase in bedridden aged people which becomes a social problem. Urgent development of a therapeutic agent for this disease is strongly desired. Disease due to a decrease in bone mass is expected to be treated by controlling bone resorption, accelerating bone formation, or improving balance between bone resorption and formation.

Osteogenesis is expected to increase by accelerating proliferation, differentiation, or activation of the cells controlling bone formation, or by controlling proliferation, differentiation, or activation of the cells involved in bone resorption. In recent years, strong interest has been directed to physiologically active proteins (cytokines) exhibiting such activities as described above, and energetic research is ongoing on this subject. The cytokines which have been reported to accelerate proliferation or differentiation of osteoblasts include the proteins of fibroblast growth factor family (FGF: Rodan S. B. et al., Endocrinology vol. 121, p 1917, 1987), insulin-like growth factor I (IGF-I: Hock J. M. et al., Endocrinology vol. 122, p 254, 1988), insulin growth factor II (IGF-II: McCarthy T. et al., Endocrinology vol. 124, p 301, 1989), Activin A (Centrella M. et al., Mol. Cell. Biol., vol. 11, p 250, 1991), transforming growth factor- β , (Noda M., The Bone, vol. 2, p 29, 1988), Vasculotropin (Varonique M. et al., Biochem. Biophys. Res. Commun., vol. 199, p 380, 1994), and the protein of heterotopic bone formation factor family (bone morphogenic protein; BMP: BMP-2; Yanaguchi A. et al., J. Cell Biol. vol. 113, p 682, 1991, OP-1; Sampath T. K. et al., J. Biol. Chem. vol. 267, p 20532, 1992, and Knutsen R. et al., Biochem. Biophys. Res. Commun. vol. 194, P 1352, 1993).

On the other hand, as the cytokines which suppress differentiation and/or maturation of osteoclasts, transforming growth factor- β (Chenu C. et al., Proc. Natl. Acad. Sci. USA, vol. 85, p 5683, 1988), interleukin-4 (Kasano K. et al., Bone-Miner., vol. 21, p 179, 1993), and the like have been reported. Further, as the cytokines which suppress bone resorption by osteoclast, calcitonin (Bone-Miner., vol. 17, p 347, 1992), macrophage colony stimulating factor (Hattersley G. et al., J. Cell. Physiol. vol. 137, p 199, 1988), interleukin-4 (Watanabe, K. et al., Biochem. Biophys. Res. Commun. vol. 172, P 1035, 1990), and interferon- γ (Gowen M. et al., J. Bone Miner. Res., vol. 1, p 46.9, 1986) have been reported.

These cytokines are expected to be used as agents for treating diseases accompanying bone loss by accelerating bone formation or suppressing of bone resorption. Clinical tests are being undertaken to verify the effect of improving bone metabolism of some cytokines such as insulin-like growth factor-I and the heterotopic bone formation factor family. In addition, calcitonin is already commercially available as a therapeutic agent for osteoporosis and a pain relief agent. At present, drugs for clinically treating bone diseases or shortening the period of treatment of bone diseases include activated vitamin D₃, calcitonin and its derivatives, and hormone preparations such as estradiol agent, ipriflavon or calcium preparations. These agents are not necessarily satisfactory in terms of the efficacy and therapeutic results. Development of a novel therapeutic agent which can be used in place of these agents is strongly desired.

In view of this situation, the present inventors have undertaken extensive studies. As a result, the present inventors had found protein OCIF exhibiting an osteoclastogenesis-inhibitory activity in a culture broth of human embryonic lung fibroblast IMR-90 (ATCC Deposition No. CCL186), and filed a patent application (PCT/JP96/00374). The present inventors have conducted further studies relating to the origin of this protein OCIF exhibiting the osteoclastogenesis-inhibitory activity. The studies have matured into determination of the sequence of a genomic DNA encoding the human origin OCIF. Accordingly, an object of the present invention is to provide a genomic DNA encoding protein OCIF exhibiting osteoclastogenesis-inhibitory activity and a process for preparing this protein by a genetic engineering technique using the genomic DNA.

DISCLOSURE OF THE INVENTION

Specifically, the present invention relates to a genomic DNA encoding protein OCIF exhibiting osteoclastogenesis-inhibitory activity and a process for preparing this protein by a genetic engineering technique using the genomic DNA.
 5 The DNA of the present invention includes the nucleotide sequences No. 1 and No. 2 in the Sequence Table attached hereto.

Moreover, the present invention relates to a process for preparing a protein, comprising inserting a DNA including the nucleotide sequences of the sequences No. 1 and No. 2 in the Sequence Table into an expression vector, producing a vector capable of expressing a protein having the following physicochemical characteristics and exhibiting the activity
 10 of inhibiting differentiation and/or maturation of osteoclasts, and producing this protein by a genetic engineering technique,

(a) molecular weight (SDS-PAGE):

- 15 (i) Under reducing conditions: about 60 kD,
 (ii) Under non-reducing conditions: about 60 kD and about 120 kD;

(b) amino acid sequence:

includes an amino acid sequence of the Sequence ID No. 3 of the Sequence Table,

20 (c) affinity:

exhibits affinity to a cation exchanger and heparin, and

(d) thermal stability:

- 25 (i) the osteoclast differentiation and/or maturation inhibitory activity is reduced when treated with heat at 70°C for 10 minutes or at 56°C for 30 minutes,

- (ii) the osteoclast differentiation and/or maturation inhibitory activity is lost when treated with heat at 90°C for 10 minutes.

The protein obtained by expressing the gene of the present invention exhibits an osteoclastogenesis-inhibitory
 30 activity. This protein is effective as an agent for the treatment and improvement of diseases involving decrease in the amount of bone such as osteoporosis, diseases relating to bone metabolism abnormality such as rheumatism, degenerative joint disease, or multiple myeloma, and is useful as an antigen to establish an immunological diagnosis of such diseases.

35 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a result of Western Blotting analysis of the protein obtained by causing genomic DNA of the present invention to express a protein in Example 4 (iii), wherein lane 1 indicates a marker, lane 2 indicates the culture broth of COS7 cells in which a vector pWESRαOCIF (Example 4 (iii)) has been transfected, and lane 3 is the culture broth of
 40 COS7 cell in which a vector pWESRα(control) has been transfected.

BEST MODE FOR CARRYING OUT THE INVENTION

The genomic DNA encoding the protein OCIF which exhibits osteoclastogenesis-inhibitory activity in the present
 45 invention can be obtained by preparing a cosmid library using a human placenta genomic DNA and a cosmid vector and by screening this library using DNA fragments which are prepared based on the OCIF cDNA as a probe. The thus-obtained genomic DNA is inserted into a suitable expression vector to prepare an OCIF expression cosmid. A recombinant type OCIF can be obtained by transfecting the genomic DNA into a host organism such as various types of cells or microorganism strains and causing the DNA to express a protein by a conventional method. The resultant protein
 50 exhibiting osteoclastogenesis-inhibitory activity (an osteoclastogenesis-inhibitory factor) is useful as an agent for the treatment and improvement of diseases involving a decrease in bone mass such as osteoporosis and other diseases relating to bone metabolism abnormality and also as an antigen to prepare antibodies for establishing immunological diagnosis of such diseases. The protein of the present invention can be prepared as a drug composition for oral or non-oral administration. Specifically, the drug composition of the present invention containing the protein which is an osteo-
 55 clastogenesis-inhibitory factor as an active ingredient can be safely administered to humans and animals. As the form of drug composition, a composition for injection, composition for intravenous drip, suppository, nasal agent, sublingual agent, percutaneous absorption agent, and the like are given. In the case of the composition for injection, such a composition is a mixture of a pharmacologically effective amount of osteoclastogenesis-inhibitory factor of the present

invention and a pharmaceutically acceptable carrier. The composition may further comprise amino acids, saccharides, cellulose derivatives, and other excipients and/or activation agents, including other organic compounds and inorganic compounds which are commonly added to a composition for injection. When an injection preparation is prepared using the osteoclastogenesis-inhibitory factor of the present invention and these excipients and activation agents, a pH adjuster, buffering agent, stabilizer, solubilizing agent, and the like may be added if necessary to prepare various types of injection agents.

The present invention will now be described in more detail by way of examples which are given for the purpose of illustration and not intended to be limiting of the present invention.

Example 1

(Preparation of a cosmid library)

A cosmid library was prepared using human placenta genomic DNA (Clontech; Cat. No. 6550-2) and pWE15 cosmid vector (Stratagene). The experiment was carried out following principally the protocol attached to the pWE15 cosmid vector kit of Stratagene Company, provided Molecular Cloning: A Laboratory Manual (Cold Spring Harbor Laboratory (1989)) was referred to for common procedures for handling DNA, E. coli, and phage.

(i) Preparation of restrictive enzymolysate of human-genomic DNA

Human placenta genomic DNA dissolved in 750 μ l of a solution containing 10 mM Tris-HCl, 10 mM MgCl₂, and 100 mM NaCl was added to four 1.5 ml Eppendorf tubes (tube A, B, C, and D) in the amount of 100 μ g each. Restriction enzyme MboI was added to these tubes in the amounts of 0.2 unit for tube A, 0.4 unit for tube B, 0.6 unit for tube C, and 0.8 unit for tube D, and DNA was digested for 1 hour. Then, EDTA in the amount to make a 20 mM concentration was added to each tube to terminate the reaction, followed by extraction with phenol/chloroform (1:1). A two-fold amount of ethanol was added to the aqueous layer to precipitate DNA. DNA was collected by centrifugation, washed with 70% ethanol, and DNA in each tube was dissolved in 100 μ l of TE (10 mM HCl (pH 8.0) + 1 mM EDTA buffer solution, hereinafter called TE). DNA in four tubes was combined in one tube and incubated for 10 minutes at 68°C. After cooling to room temperature, the mixture was overlayed onto a 10%-40 % linear sucrose gradient which was prepared in a buffer containing 20 mM Tris-HCl (pH 8.0), 5 mM EDTA, and 1 mM NaCl in a centrifugal tube (38 ml). The tube was centrifuged at 26,000 rpm for 24 hours at 20°C using a rotor SRP28SA manufactured by Hitachi, Ltd. and 0.4 ml fractions of the sucrose gradient was collected using a fraction collector. A portion of each fraction was subjected to 0.4% agarose electrophoresis to confirm the size of DNA. Fractions containing DNA with a length of 30 kb (kilo base pair) to 40 kb were thus combined. The DNA solution was diluted with TE to make a sucrose concentration to 10% or less and 2.5-fold volumes of ethanol was added to precipitate DNA. DNA was dissolved in TE and stored at 4°C.

(ii) Preparation of cosmid vector

The pWE15 cosmid vector obtained from Stratagene Company was completely digested with restriction enzyme BamHI according to the protocol attached to the cosmid vector kit. DNA collected by ethanol precipitation was dissolved in TE to a concentration of 1 mg/ml. Phosphoric acid at the 5'-end of this DNA was removed using calf small intestine alkaline phosphatase, and DNA was collected by phenol extraction and ethanol precipitation. The DNA was dissolved in TE to a concentration of 1 mg/ml.

(iii) Ligation of genomic DNA to vector and in vitro packaging

1.5 micrograms of genomic DNA fractionated according to size and 3 μ g of pWE15 cosmid vector which was digested with restriction enzyme BamHI were ligated in 20 μ l of a reaction solution using Ready-To-Go T4DNA ligase of Pharmacia Company. The ligated DNA was packaged in vitro using Gigapack™ II packaging extract (Stratagene) according to the protocol. After the packaging reaction, a portion of the reaction mixture was diluted stepwise with an SM buffer solution and mixed with E. coli XL1-Blue MR (Stratagene) which was suspended in 10 mM MgCl₂ to cause phage to infect, and plated onto LB agar plates containing 50 μ g/ml of ampicillin. The number of colonies produced was counted. The number of colonies per 1 μ l of packaging reaction was calculated based on this result.

(iv) Preparation of a cosmid library

The packaging reaction solution thus prepared was mixed with E. coli XL1-Blue MR and the mixture was plated onto agarose plates containing ampicillin so as to produce 50,000 colonies per agarose plate having a 15 cm of diam-

eter. After incubating the plate overnight at 37°C, an LB culture medium was added in the amount of 3 ml per plate to suspend and collect colonies of *E. coli*. Each agarose plate was again washed with 3 ml of the LB culture medium and the washing was combined with the original suspension of *E. coli*. The *E. coli* collected from all agarose plates was placed in a centrifugal tube, glycerol was added to a concentration of 20%, and ampicillin was further added to make a final concentration of 50 µg/ml. A portion of the *E. coli* suspension was removed and the remainder was stored at -80°C. The removed *E. coli* was diluted stepwise and plated onto an agar plates to count the number of colonies per 1 ml of suspension.

Example 2

(Screening of cosmid library and purification of colony)

A nitrocellulose filter (Millipore) with a diameter of 14.2 cm was placed on each LB agarose plate with a diameter of 15 cm which contained 50 µg/ml of ampicillin. The cosmid library was plated onto the plates so as to produce 50,000 colonies of *E. coli* per plate, followed by incubation overnight at 37°C. *E. coli* on the nitrocellulose filter was transferred to another nitrocellulose filter according to a conventional method to obtain two replica filters. According to the protocol attached to the cosmid vector kit, cosmid DNA in the *E. coli* on the replica filters was denatured with an alkali, neutralized, and immobilized on the nitrocellulose filter using a Stratalinker (Stratagene). The filters were heated for two hours at 80°C in a vacuum oven. The nitrocellulose filters thus obtained were hybridized using two kinds of DNA produced, respectively, from 5'-end and 3'-end of human OCIF cDNA as probes. Namely, a plasmid was purified from *E. coli* pKB/OIF10 (deposited at The Ministry of International Trade and Industry, the Agency of Industrial Science and Technology, Biotechnology Laboratory, Deposition No. FERM BP-5267) containing OCIF cDNA. The plasmid containing OCIF cDNA was digested with restriction enzymes KpnI and EcoRI. Fragments thus obtained was separated using agarose gel electrophoresis. KpnI/EcoRI fragment with a length of 0.2 kb was purified using a QIAEX II gel extraction kit (Qiagen). This DNA was labeled with ³²P using the Megaprime DNA Labeling System (Amasham) (5'-DNA probe). Apart from this, a BamHI/EcoRV fragment with a length of 0.2 kb which was produced from the above plasmid by digestion with restriction enzymes BamHI and EcoRV was purified and labeled with ³²P (3'-DNA probe). One of the replica filters described above was hybridized with the 5'-DNA probe and the other with the 3'-DNA probe. Hybridization and washing of the filters were carried out according to the protocol attached to the cosmid vector kit. Autoradiography detected several positive signals with each probe. One colony which gave positive signals with both probe was identified. The colony on the agar plate, which corresponding to the signal on the autoradiogram was isolated and purified. A cosmid was prepared from the purified colony by a conventional method. This cosmid was named pWEOCIF. The size of human genomic DNA contained in this cosmid was about 38 kb.

Example 3

(Determination of the nucleotide sequence of human OCIF genomic DNA)

(i) Subcloning of OCIF genomic DNA

Cosmid pWEOCIF was digested with restriction enzyme EcoRI. After the separation of the DNA fragments thus produced by electrophoresis using a 0.7% agarose gel, the DNA fragments were transferred to a nylon membrane (Hybond -N, Amasham) by the Southern blot technique and immobilized on the nylon membrane using Stratalinker (Stratagene). On the other hand, plasmid pBKOCIF was digested with restriction enzyme EcoRI and a 1.6 kb fragment containing human OCIF cDNA was isolated by agarose gel electrophoresis. The fragment was labeled with ³²P using the Megaprime DNA labeling system (Amasham).

Hybridization of the nylon membranes described above with the ³²P-labeled 1.6-kb OCIF cDNA was performed according to a conventional method detected that DNA fragments with a size of 6 kb, 4 kb, 3.6 kb, and 2.6 kb. These fragments hybridized with the human OCIF cDNA were isolated using agarose gel electrophoresis and individually subcloned into an EcoRI site of pBluescript II SK + vector (Stratagene) by a conventional method. The resulting plasmids were respectively named pBSE 6, pBSE 4, pBSE 3.6, and pBSE 2.6.

(ii) Determination of the nucleotide sequence

The nucleotide sequence of human OCIF genomic DNA which was subcloned into the plasmid was determined using the ABI Dideoxy Terminator Cycle Sequencing Ready Reaction kit (Perkin Elmer) and the 373 Sequencing System (Applied Biosystems). The primer used for the determination of the nucleotide sequence was synthesized based on the nucleotide sequence of human OCIF cDNA (Sequence ID No. 4 in the Sequence Table). The nucleotide

sequences thus determined are given as the Sequences No. 1 and No. 2 in the Sequence Table. The Sequence ID No. 1 includes the first exon of the OCIF gene and the Sequence ID No. 2 includes the second, third, fourth, and fifth exons. A stretch of about 17 kb is present between the first and second exons.

5 Example 4

(Production of recombinant OCIF using COS-7 cells)

(i) Preparation of OCIF genomic DNA expression cosmid

10

To express OCIF genomic DNA in animal cells, an expression unit of expression plasmid pcDL-SR α 296 (Molecular and Cellular Biology, vol. 8, P466-472, 1988) was inserted into cosmid vector pWE15 (Stratagene). First of all, the expression plasmid pcDL-SR α 296 was digested with a restriction enzyme Sal I to cut out expression unit with a length of about 1.7 kb which includes an SR α promotor, SV40 later splice signal, poly (A) addition signal, and so on. The digestion products were separated by agarose electrophoresis and the 1.7-kb fragment was purified using the QIAEX II gel extraction kit (Qiagen). On the other hand, cosmid vector pWE15 was digested with a restriction enzyme EcoRI and fragments were separated using agarose gel electrophoresis. pWE15 DNA of 8.2 kb long was purified using the QIAEX II gel extraction kit (Qiagen). The ends of these two DNA fragments were bluntended using a DNA blunting kit (Takara Shuzo), ligated using a DNA ligation kit (Takara Shuzo), and transferred into *E. coli* DH5 α (Gibco BRL). The resultant transformant was grown and the expression cosmid pWESR α containing an expression unit was purified using a Qiagen column (Qiagen).

The cosmid pWE OCIF containing the OCIF genomic DNA with a length of about 38 kb obtained in (i) above was digested with a restriction enzyme NotI to cut out the OCIF genomic DNA of about 38 kb. After separation by agarose gel electrophoresis, the DNA was purified using the QIAEX II gel extraction kit (Qiagen). On the other hand, the expression cosmid pWESR α was digested with a restriction enzyme EcoRI and the digestion product was extracted with phenol and chloroform, ethanol-precipitated, and dissolved in TE.

pWESR α digested with a restriction enzyme EcoRI and an EcoRI-XmnI-NotI adapter (#1105, #1156 New England Biolaboratory Co.) were ligated using T4 DNA ligase (Takara Shuzo Co., Ltd.). After removal of the free adapter by agarose gel electrophoresis, the product was purified using QIAEX gel extraction kit (Qiagen). The OCIF genomic DNA with a length of about 37 kb which was derived from the digestion with restriction enzyme NotI and the pWESR α to which the adapter was attached were ligated using T4 DNA ligase (Takara Shuzo). The DNA was packaged in vitro using the Gigapack packaging extract (Stratagene) and infected with *E. coli* XL1-Blue MR (Stratagene). The resultant transformant was grown and the expression cosmid pWESR α OCIF which contained OCIF genomic DNA was inserted was purified using a Qiagen column (Qiagen). The OCIF expression cosmid pWESR α OCIF was ethanol-precipitated and dissolved in sterile distilled water and used in the following analysis.

(ii) Transient expression of OCIF genomic DNA and measurement of OCIF activity

A recombinant OCIF was expressed as described below using the OCIF expression cosmid pWESR α OCIF obtained in (i) above and its activity was measured. COS-7 (8×10^5 cells/well) cells (Riken Cell Bank, RCB0539) were planted in a 6-well plate using DMEM culture medium (Gibco BRL) containing 10% fetal bovine serum (Gibco BRL). On the following day, the culture medium was removed and cells were washed with serum-free DMEM culture medium. The OCIF expression cosmid pWESR α OCIF which had been diluted with OPTI-MEM culture medium (Gibco BRL) was mixed with lipopectamine and the mixture was added to the cells in each well according to the attached protocol. The expression cosmid pWESR α was added to the cells in the same manner as a control. The amount of the cosmid DNA and Lipopectamine was respectively 3 μ g and 12 μ l. After 24 hours, the culture medium was removed and 1.5 ml of fresh EX-CELL 301 culture medium (JRH Bioscience) was added to each well. The culture medium was recovered after 48 hours and used as a sample for the measurement of OCIF activity. The measurement of OCIF activity was carried out according to the method described by Kumegawa, M. et al. (Protein, Nucleic Acid, and Enzyme, Vol. 34, p 999 (1989)) and the method of TAKAHASHI, N. et al. (Endocrinology vol. 122, p 1373 (1988)). The osteoclast formation in the presence of activated vitamin D₃ from bone marrow cells isolated from mice aged about 17 days was evaluated by the induction of tartaric acid resistant acidic phosphatase activity. The inhibition of the acid phosphatase was measured and used as the activity of the protein which possesses osteoclastogenesis-inhibitory activity (OCIF). Namely, 100 μ l/well of a OCIF sample which was diluted with α -MEM culture medium (Gibco BRL) containing 2×10^{-8} M activated vitamin D₃ and 10% fetal bovine serum was added to each well of a 96 well micro plate. Then, 3×10^5 bone marrow cells isolated from mice (about 17-days old) suspended in 100 μ l of α -MEM culture medium containing 10% fetal bovine serum were added to each well of the 96 well micro plate and cultured for a week at 37°C and 100% humidity under 5% CO₂ atmosphere. On days 3 and 5, 160 μ l of the conditioned medium was removed from each well, and 160 μ l of a sam-

ple which was diluted with α -MEM culture medium containing 1×10^{-8} M activated vitamin D₃ and 10% fetal bovine serum was added. After 7 days from the start of culturing, the cells were washed with a phosphate buffered saline and fixed with a ethanol/acetone (1:1) solution for one minute at room temperature. The osteoclast formation was detected by staining the cells using an acidic phosphatase activity measurement kit (Acid Phosphatase, Leucocyte, Cat.No. 387-A, Sigma Company). A decrease in the number of cells positive to acidic phosphatase activity in the presence of tartaric acid was taken as the OCIF activity. The results are shown in Table 1, which indicates that the conditioned medium exhibits the similar activity to natural type OCIF obtained from the IMR-90 culture medium and recombinant OCIF produced by CHO cells.

TABLE 1

Activity of OCIF expressed by COS-7 cells in the conditioned medium						
Dilution	1/10	1/20	1/40	1/80	1/160	1/320
OCIF genomic DNA introduced	++	++	++	++	+	-
Vector introduced	-	-	-	-	-	-
Untreated	-	-	-	-	-	-
"++" indicates an activity inhibiting 80% or more of osteoclast formation, "+" indicates an activity inhibiting 30-80% of osteoclast formation, and "-" indicates that no inhibition of osteoclast formation is observed.						

(iii) Identification of the product by Western Blotting

A buffer solution (10 μ l) for SDS-PAGE (0.5 M Tris-HCl, 20% glycerol, 4% SDS, 20 μ g/ml bromophenol blue, pH 6.8) was added to 10 μ l of the sample for the measurement of OCIF activity prepared in (ii) above. After boiling for 3 minutes at 100°C, the mixture was subjected to 10% SDS polyacrylamide electrophoresis under non-reducing conditions. The proteins were transferred from the gel to a PVDF membrane (ProBlott, Perkin Elmer) using semi-dry blotting apparatus (Biorad). The membrane was blocked and incubated for 2 hours at 37°C together with a horseradish peroxidase-labeled anti-OCIF antibody obtained by labeling the previously obtained OCIF protein with horseradish peroxidase according to a conventional method. After washing, the protein which has bound the anti-OCIF antibody was detected using the ECL system (Amasham). As shown in Figure 1, two bands, one with a molecular weight of about 120 kilo dalton and the other 60 kilo dalton, were detected in the supernatant obtained from the culture broth of COS-7 cells in which pWESR α OCIF was transfected. On the other hand, these two bands with a molecular weight of about 120 kilo dalton and 60 kilo dalton were not detected in the supernatant obtained from the culture broth of COS-7 cells in which pWESR α vector was transfected, confirming that the protein obtained was OCIF.

INDUSTRIAL APPLICABILITY

The present invention provides a genomic DNA encoding a protein OCIF which possesses an osteoclastogenesis-inhibitory activity and a process for preparing this protein by a genetic engineering technique using the genomic DNA. The protein obtained by expressing the gene of the present invention exhibits an osteoclastogenesis-inhibitory activity and is useful as an agent for the treatment and improvement of diseases involving a decrease in the amount of bone such as osteoporosis, other diseases resulting from bone metabolism abnormality such as rheumatism or degenerative joint disease, and multiple myeloma. The protein is further useful as an antigen to establish antibodies useful for an immunological diagnosis of such diseases.

NOTE ON MICROORGANISM

Depositing Organization:

The Ministry of International Trade and Industry, National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology

Address: 1-3, Higashi-1-Chome, Tsukuba-shi, Ibaraki-ken, Japan

Date of Deposition: June 21, 1995 (originally deposited on June 21, 1995 and transferred to the international deposition according to the Budapest Treaty on October 25, 1995)

Accession No. FERM BP-5267

TABLE OF SEQUENCES

Sequence number: 1

Length of sequence: 1316

Sequence Type: nucleic acid

Strandedness: double

Topology: linear

Molecular type: genomic DNA (human OCIF genomic DNA-1)

Sequence:

```

CTGGAGACAT ATAAGTTGAA CACTTGGCCC TGATGGGGAA GCAGCTCTGC AGGGACTTTT  60
TCAGCCATCT GTAAACAATT TCAGTGGCAA CCCGCCAACT GTAATCCATG AATGGGACCA 120
CACTTTACAA GTCATCAAGT CTAAGTTCTA GACCAGGGAA TTAATCGGGG AGACAGCGAA 180
CCCTAGAGCA AAGTGCCAAA CTTCTGTGGA TAGCTTGAGG CTAGTGGAAA GACCTCGAGG 240
AGGCTACTCC AGAAGTTCAG GCGGTAGGAA GCTCCGATAC CAATAGCCCT TTGATGATGG 300
TGGGGTTGGT GAAGGGAACA GTGCTCCGCA AGGTTATCCC TGCCCCAGGC AGTCCAATTT 360
TCACTCTGCA GATTCTCTCT GGCTCTAACT ACCCCAGATA ACAAGGAGTG AATGCAGAAT 420
AGCAGCGGCT TTAGGGCCAA TCAGACATTA GTTAGAAAAA TTCCTACTAC ATGGTTTATG 480
TAAAGTTGAA GATGAATCAT TCGGAAGTCC CCGAAAAGGG CTCAGACAAT GCCATGCATA 540
AAGAGGGGCC CTGTAATTTG AGGTTTCAGA ACCCGAAGTG AAGGGGTCAG GCAGCCGGGT 600
ACGGCGGAAA CTCACAGCTT TCGCCAGCG AGAGGACAAA GGTCTGGGAC AACTCCAAC 660
TGGCTCCGGA TCTTGGCTGG ATCGGACTCT CAGGGTGGAG GAGACACAAG CACAGCAGCT 720
GCCCAGCGTG TCCCAGCCC TCCACCGCT GTTCCCGCT GCCAGGAGGC TGGCCGCTGG 780
CGGGAAGGGG CCGGGAACCC TCAGAGCCCC GCGGAGACAG CAGCCGCTT GTTCTCAGC 840
CCGCTGGCTT TTTTTCCTCC TGCTCTCCCA GGGGACAGAC ACCACCGCCC CACCCCTCAC 900
GCCCCACCTC CCTGGGGGAT CTTTCCGCC CCAGCCCTGA AAGCGTTAAT CCTGGAGCTT 960
TCTGCACACC CCCCAGCCG TCCGCCCCAA GCTTCTAAA AAAGAAAGGT GCAAAGTTTG 1020
GTCCAGGATA GAAAAATCAC TGATCAAAGG CAGGCGATAC TTCCTGTTGC CGGGACGCTA 1080
TATATAACGT GATGAGCGCA CGGCTGCGG AGACGCACCG GAGCGCTCGC CCAGCCGCGC 1140

```

5 CCTCCAAGCC CCTGAGGTTT CCGGGGACCA CA ATG AAC AAG TTG CTG TGC TGC 1193
 Met Asn Lys Leu Leu Cys Cys
 -20 -15

10 GCG CTC GTG GTAAGTCCCT GGGCCAGCCG ACGGGTGCCC GCGGCCTGGG 1242
 Ala Leu Val

15 GAGGCTGCTG CCACCTGGTC TCCCAACCTC CCAGCGGACC GGCGGGGAGA AGGCTCCACT 1302
 CGCTCCCTCC CAGG 1316

20 Sequence number: 2
 Length of sequence: 9898
 25 Sequence Type: nucleic acid
 Strandedness: double
 Topology: linear
 30 Molecular type: genomic DNA (human OCIF genomic DNA-2)

35 Sequence:
 GCTTACTTTG TGCCAAATCT CATTAGGCTT AAGGTAATAC AGGACTTTGA GTCAAATGAT 60
 ACTGTTGCAC ATAAGAACAA ACCTATTTTC ATGCTAAGAT GATGCCACTG TGTTCCTTTC 120
 40 TCCTTCTAG TTT CTG GAC ATC TCC ATT AAG TGG ACC ACC CAG GAA ACG TTT 171
 Phe Leu Asp Ile Ser Ile Lys Trp Thr Thr Gln Glu Thr Phe
 45 -10 -5 1

CCT CCA AAG TAC CTT CAT TAT GAC GAA GAA ACC TCT CAT CAG CTG TTG 219
 50 Pro Pro Lys Tyr Leu His Tyr Asp Glu Glu Thr Ser His Gln Leu Leu
 5 10 15

55

5 TGT GAC AAA TGT CCT CCT GGT ACC TAC CTA AAA CAA CAC TGT ACA GCA 267
 Cys Asp Lys Cys Pro Pro Gly Thr Tyr Leu Lys Gln His Cys Thr Ala
 20 25 30 35

10 AAG TGG AAG ACC GTG TGC GCC CCT TGC CCT GAC CAC TAC TAC ACA GAC 315
 Lys Trp Lys Thr Val Cys Ala Pro Cys Pro Asp His Tyr Tyr Thr Asp
 15 40 45 50

20 AGC TGG CAC ACC AGT GAC GAG TGT CTA TAC TGC AGC CCC GTG TGC AAG 363
 Ser Trp His Thr Ser Asp Glu Cys Leu Tyr Cys Ser Pro Val Cys Lys
 25 55 60 65

30 GAG CTG CAG TAC GTC AAG CAG GAG TGC AAT CGC ACC CAC AAC CGC GTG 411
 Glu Leu Gln Tyr Val Lys Gln Glu Cys Asn Arg Thr His Asn Arg Val
 70 75 80

35 TGC GAA TGC AAG GAA GGG CGC TAC CTT GAG ATA GAG TTC TGC TTG AAA 459
 Cys Glu Cys Lys Glu Gly Arg Tyr Leu Glu Ile Glu Phe Cys Leu Lys
 40 85 90 95

45 CAT AGG AGC TGC CCT CCT CGA TTT CGA GTG GTG CAA GCT G GTACGTGTCA 509
 His Arg Ser Cys Pro Pro Gly Phe Gly Val Val Gln Ala
 100 105 110

50 ATGTGCAGCA AAATTAATTA GGATCATGCA AAGTCAGATA GTTGTGACAG TTTAGGAGAA 569

55

CACTTTTGT CTGATGACAT TATAGGATAG CAAATTGCAA AGGTAATGAA ACCTGCCAGG 629
 TAGGTACTAT GTGTCTGGAG TGCTTCCAAA GGACCATTCG TCAGAGGAAT ACTTTGCCAC 689
 5 TACAGGGCAA TTTAATGACA AATCTCAAAT GCAGCAAATT ATTCTCTCAT GAGATGCATG 749
 ATCGTTTTTT TTTTTTTTTT TAAAGAAACA AACTCAAGTT GCACTATTGA TAGTTGATCT 809
 10 ATACCTCTAT ATTTCACTTC AGCATGGACA CCTTCAAACG GCAGCACTTT TTGACAAACA 869
 TCAGAAATGT TAATTATAC CAAGAGAGTA ATTATGCTCA TATTAATGAG ACTCTGGAGT 929
 GCTAACAATA AGCAGTTATA ATTAATTATG TAAAAAATGA GAATGGTGAG GGAATTGCA 989
 15 TTTCATTATT AAAACAAGG CTAGTTCTTC CTTAGCATG GGAGCTGAGT GTTGGGAGG 1049
 GTAAGGACTA TAGCAGAATC TCTTCAATGA GCTTATTCTT TATCTTAGAC AAAACAGATT 1109
 20 GTCAAGCCAA GAGCAAGCAC TTGCTATAA ACCAAGTGCT TTCTCTTTTG CATTTTGAAC 1169
 AGCATTGGTC AGGGCTCATG TGTATTGAAT CTTTAAACC AGTAACCCAC GTTTTTTTTC 1229
 TGCCACATTT GCGAAGCTTC AGTGCAGCCT ATAACTTTC ATAGCTTGAG AAAATTAAGA 1289
 25 GTATCCACTT ACTTAGATGG AAGAAGTAAT CAGTATAGAT TCTGATGACT CAGTTTGAAG 1349
 CAGTGTCTCT CAACTGAAGC CCTGCTGATA TTTAAGAAA TATCTGGATT CCTAGGCTGG 1409
 30 ACTCCTTTTT GTGGGCAGCT GTCCTGCGCA TTGTAGAATT TTGGCAGCAC CCCTGGAATC 1469
 TAGCCACTAG ATACCAATAG CAGTCCTTCC CCCATGTGAC AGCCAAAAAT GTCTTCAGAC 1529
 ACTGTCAAAT GTCGCCAGGT GGCAAAATCA CTCCTGGTTG AGAACAGGGT CATCAATGCT 1589
 35 AAGTATCTGT AACTATTTTA ACTCTCAAAA CTTGTGATAT ACAAAGTCTA AATTATTAGA 1649
 CGACCAATAC TTTAGGTTTA AAGGCATACA AATGAAACAT TCAAAAATCA AAATCTATTC 1709
 40 TGTTTCTCAA ATAGTGAATC TTATAAAATT AATCACAGAA GATGCAAATT GCATCAGACT 1769
 CCCTTAAAT TCCTCTTCGT ATGAGTATTT GAGGGAGGAA TTGGTGATAG TTCCTACTTT 1829
 CTATTGGATG GTACTTTGAG ACTCAAAAGC TAAGCTAAGT TGTGTGTGTG TCAGGGTGCG 1889
 45 GGGTGTGGAA TCCCATCAGA TAAAGCAAA TCCATGTAAT TCATTCAGTA AGTTGTATAT 1949
 GTAGAAAAAT GAAAAGTGGG CTATGCAGCT TGGAACTAG AGAATTTTGA AAAATAATGG 2009
 50 AAATCACAAG GATCTTTCTT AAATAAGTAA GAAAATCTGT TTGTAGAATG AAGCAAGCAG 2069
 GCAGCCAGAA GACTCAGAAC AAAAGTACAC ATTTTACTCT GTGTACTCTG GCAGCACAGT 2129

55

GGGATTTATT TACCTCTCCC TCCCTAAAAA CCCACACAGC GGTTCCTCTT GGGAAATAAG 2189
 5 AGGTTTCCAG CCCAAAGAGA AGGAAAGACT ATGTGGTGTT ACTCTAAAAA GTATTTAATA 2249
 ACCGTTTTGT TGTTCCTGTT GCTGTTTTGA AATCAGATTG TCTCCTCTCC ATATTTTATT 2309
 TACTTCATTC TGTAAATTC TGTGGAATTA CTTAGAGCAA GCATGGTGAA TTCTCAACTG 2369
 10 TAAAGCCAAA TTTCTCCATC ATTATAATTT CACATTTTGC CTGGCAGGTT ATAATTTTAA 2429
 TATTTCCACT GATAGTAATA AGGTAAATC ATTACTTAGA TGGATAGATC TTTTTCATAA 2489
 AAAGTACCAT CAGTTATAGA GGAAGTCAT GTTCATGTTT AGGAAGGTCA TTAGATAAAG 2549
 15 CTTCTGAATA TATTATGAAA CATTAGTTCT GTCATTCTTA GATTCTTTTT GTTAAATAAC 2609
 TTTAAAAGCT AACTTACCTA AAAGAAATAT CTGACACATA TGAAGTCTC ATTAGGATGC 2669
 20 AGGAGAAGAC CCAAGCCACA GATATGTATC TGAAGAATGA ACAAGATTCT TAGGCCCCGC 2729
 ACGGTGGCTC ACATCTGTAA TCTCAAGAGT TTGAGAGGTC AAGGCGGGCA GATCACCTGA 2789
 GGTCAGGAGT TCAAGACCAG CCTGGCCAAC ATGATGAAAC CCTGCCTCTA CTAAAAATAC 2849
 25 AAAAATTAGC AGGGCATGCT GGTGCATGCC TGCAACCCTA GCTACTCAGG AGGCTGAGAC 2909
 AGGAGAATCT CTTGAACCTT CGAGCGGAG GTTGTGGTGA GCTGAGATCC CTCTACTGCA 2969
 30 CTCCAGCCTG GGTGACAGAG ATGAGACTCC GTCCCTGCCG CCGCCCCCGC CTTCCCCCCC 3029
 AAAAAGATTC TTCTTCATGC AGAACATACG GCAGTCAACA AAGGGAGACC TGGGTCCAGG 3089
 TGTCCAAGTC ACTTATTTGG AGTAAATTAG CAATGAAAGA ATGCCATGGA ATCCCTGCCC 3149
 35 AAATACCTCT GCTTATGATA TTGTAGAATT TGATATAGAG TTGTATCCCA TTTAAGGAGT 3209
 AGGATGTAGT AGGAAAGTAC TAAAAACAAA CACACAAACA GAAAACCTC TTTGCTTTGT 3269
 40 AAGGTGGTTC CTAAGATAAT GTCAGTGCAA TGCTGAAAT AATATTTAAT ATGTGAAGGT 3329
 TTTAGGCTGT GTTTTCCCTT CCTGTTCTTT TTTCTGCCA GCCCTTTGTC ATTTTGCAG 3389
 GTCAATGAAT CATGTAGAAA GAGACAGGAG ATGAAACTAG AACCAGTCCA TTTTCCCCCT 3449
 45 TTTTTATTT TCTGGTTTTG GTAAAAGATA CAATGAGGTA GGAGGTTGAG ATTTATAAAT 3509
 GAAGTTTAAT AAGTTTCTGT AGCTTTGATT TTTCTCTTC ATATTGTGA TCTGCATAA 3569
 50 GCCAGAAATG GCCTGTAAAA TCTACATATG GATATTGAAG TCTAAATCTG TTCAACTAGC 3629
 TTACTACTAGA TGGAGATATT TTCAATTTCA GATACACTGG AATGTATGAT CTAGCCATGC 3689

5 GTAATATAGT CAAGTGTGTTG AAGGTATTIA TTTTAAATAG CGTCTTTAGT TGTGGACTGG 3749
 TTCAAGTTTT TCTGCCAATG ATTTCTTCAA ATTTATCAAA TATTTTCCCA TCATGAAGTA 3809
 AAATGCCCTT GCAGTCACCC TTCCTGAAGT TTGAACGACT CTGCTGTTTT AAACAGTTTA 3869
 AGCAAATGGT ATATCATCTT CCGTTTACTA TGTAGCTTAA CTGCAGGCTT ACGCTTTTGA 3929
 10 GTCAGCGGCC AACTTTATTG CCACCTTCAA AAGTTTATTA TAATGTTGTA AATTTTACT 3989
 TCTCAAGGTT AGCATACTTA GGAGTTGCTT CACAATTAGG ATTCAGGAAA GAAAGAACTT 4049
 CAGTAGGAAC TGATTGGAAT TTAATGATGC AGCATTCAAT GGGTACTAAT TTCAAAGAAT 4109
 15 GATATTACAG CAGACACACA GCAGTTATCT TGATTTTCTA GGAATAATTG TATGAAGAAT 4169
 ATGGCTGACA ACACGGCCTT ACTGCCACTC AGCGGAGGCT GGACTAATGA ACACCCTACC 4229
 20 CTTCTTTTCTT TTCCTCTCAC ATTTTCATGAG CGTTTTGTAG GTAACGAGAA AATTGACTTG 4289
 CATTTCGATT ACAAGGAGGA GAAACTGGCA AAGGGGATGA TGGTGGAAGT TTTGTTCTGT 4349
 25 CTAATGAAGT GAAAAATGAA AATGCTAGAG TTTTGTGCAA CATAACTAGTA GCAGTAAAAA 4409
 CCAAGTGAAA AGTCTTTTCCA AACTGTGTT AAGAGGGCAT CTGCTGGGAA ACGATTTGAG 4469
 GAGAAGGTAC TAAATTGCTT GGTATTTTCC GTAG GA ACC CCA GAG CGA AAT ACA 4523

Gly Thr Pro Glu Arg Asn Thr

115

35 GTT TGC AAA AGA TGT CCA GAT GGG TTC TTC TCA AAT CAG ACG TCA TCT 4571
 Val Cys Lys Arg Cys Pro Asp Gly Phe Phe Ser Asn Glu Thr Ser Ser
 40 120 125 130 135

45 AAA GCA CCC TGT AGA AAA CAC ACA AAT TGC AGT GTC TTT GGT CTC CTG 4619
 Lys Ala Pro Cys Arg Lys His Thr Asn Cys Ser Val Phe Gly Leu Leu
 140 145 150

50 CTA ACT CAG AAA GGA AAT GCA ACA CAC GAC AAC ATA TGT TCC GGA AAC 4667
 55

Leu Thr Gln Lys Gly Asn Ala Thr His Asp Asn Ile Cys Ser Gly Asn

155

160

165

AGT GAA TCA ACT CAA AAA TGT GGA ATA G GTAATTACAT TCCAAAATAC 4715

Ser Glu Ser Thr Gln Lys Cys Gly Ile

170

175

GTCTTTGTAC GATTTTGTAG TATCATCTCT CTCTCTGAGT TGAACACAAG GCCTCCAGCC 4775

ACATTCTTGG TCAAACCTTAC ATTTTCCCTT TCTTGAATCT TAACCAGCTA AGGCTACTCT 4835

CGATGCATTA CTGCTAAAGC TACCACTCAG AATCTCTCAA AAATCATCT TCTCACAGAT 4895

AACACCTCAA AGCTTCATTT TCTCTCCTTT CACACTGAAA TCAAATCTTG CCCATAGGCA 4955

AAGGGCAGTG TCAAGTTTGC CACTGAGATG AAATTAGGAG AGTCCAAACT GTAGAATTCA 5015

CGTTGTGTGT TATTACTTTC ACGAATGTCT GTATTATTAA CTAAAGTATA TATTGGCAAC 5075

TAAGAAGCAA AGTGATATAA ACATGATGAC AAATTAGGCC AGGCATGGTG GCTTACTCCT 5135

ATAATCCCAA CATTTTGGGG GGCCAAGGTA GGCAGATCAC TTGAGGTCAG GATTTCAAGA 5195

CCAGCCTGAC CAACATGGTG AAACCTTGTC TCTACTAAAA ATACAAAAAT TAGCTGGGCA 5255

TGGTAGCAGG CACTTCTAGT ACCAGCTACT CAGGGCTGAG GCAGGAGAAT CGCTTGAACC 5315

CAGGAGATGG AGGTTGCAGT GAGCTGAGAT TGTACCACTG CACTCCAGTC TGGGCAACAG 5375

AGCAAGATTT CATCACACAC ACACACACAC ACACACACAC ACACATTAGA AATGTGTACT 5435

TGGCTTTGTT ACCTATGGTA TTAGTGCATC TATTGCATGG AACTTCCAAG CTA CTCTCGT 5495

TGTGTTAAGC TCTTCATTGG GTACAGGTCA CTAGTATTAA GTTCAGGTTA TTCGGATGCA 5555

TTCCACGGTA GTGATGACAA TTCATCAGGC TAGTGTGTGT GTTCACCTTG TCACTCCCAC 5615

CACTAGACTA ATCTCAGACC TTCACTCAA GACACATTAC ACTAAAGATG ATTTGCTTTT 5675

TTGTGTTTAA TCAAGCAATG GTATAAACCA GCTTGACTCT CCCCACACAG TTTTTCGTAC 5735

TACAAAGAAG TTTATGAAGC AGAGAAATGT GAATTGATAT ATATATGAGA TTCTAACCCA 5795

GTTCCAGCAT TGTTTCATTG TGTAATTGAA ATCATAGACA AGCCATTTTA GCCTTTCCTT 5855

TCTTATCTAA AAAAAAAAAA AAAAAAATGA AGGAAGGGGT ATTAAAAGGA GTGATCAAAT 5915
 5 TTTAACATTC TCTTTAATTA ATTCATTTTT AATTTTACTT TTTTTCATTT ATTGTGCACT 5975
 TACTATGTGG TACTGTGCTA TAGAGGCTTT AACATTTATA AAAACACTGT GAAAGTTGCT 6035
 TCAGATGAAT ATAGGTAGTA GAACGGCAGA ACTAGTATTC AAAGCCAGGT CTGATGAATC 6095
 10 CAAAAACAAA CACCCATTAC TCCCATTTTC TGGGACATAC TTACTCTACC CAGATGCTCT 6155
 GGGCTTTGTA ATGCCTATGT AAATAACATA GTTTTATGTT TGGTTATTTT CCTATGTAAT 6215
 GTCTACTTAT ATATCTGTAT CTATCTCTTG CTTTGTITCC AAAGGTAAAC TATGTGTCTA 6275
 15 AATGTGGGCA AAAAATAACA CACTATTECA AATTACTGTT CAAATTCCTT TAAGTCAGTG 6335
 ATAATTATTT GTTTTGACAT TAATCATGAA GTTCCCTGTG GGTACTAGGT AAACCTTTAA 6395
 20 TAGAATGTTA ATGTTTGTAT TCATTATAAG AATTTTTGGC TGTTACTTAT TTACAACAAT 6455
 ATTTCACTCT AATTAGACAT TTACTAAACT TTCTCTTGAA AACAATGCCC AAAAAAGAAC 6515
 ATTAGAAGAC ACCTAAGCTC AGTTGGTCTC TGCCACTAAG ACCAGCCAAC AGAAGCTTGA 6575
 25 TTTTATTCAA ACTTTGCATT TTAGCATATT TTATCTTGGA AAATTCAATT GTGTTGGTTT 6635
 TTTGTTTTTG TTTGTATTGA ATAGACTCTC AGAAATCCAA TTGTTGAGTA AATCTTCTGG 6695
 30 GTTTTCTAAC CTTTCTTTAG AT GTT ACC CTG TGT GAG GAG GCA TTC TTC AGG 6747

Asp Val Thr Leu Cys Glu Glu Ala Phe Phe Arg

180

185

35

TTT GCT GTT CCT ACA AAG TTT ACG CCT AAC TGG CTT AGT GTC TTG GTA 6795
 40 Phe Ala Val Pro Thr Lys Phe Thr Pro Asn Trp Leu Ser Val Leu Val

190

195

200

45

GAC AAT TTG CCT GGC ACC AAA GTA AAC GCA GAG AGT GTA GAG AGG ATA 6843
 Asp Asn Leu Pro Gly Thr Lys Val Asn Ala Glu Ser Val Glu Arg Ile

50

205

210

215

55

AAA CGG CAA CAC AGC TCA CAA GAA CAG ACT TTC CAG CTG CTG AAG TTA 6891

Lys Arg Gln His Ser Ser Gln Glu Gln Thr Phe Gln Leu Leu Lys Leu

220

225

230

235

TGG AAA CAT CAA AAC AAA GAC CAA GAT ATA GTC AAG AAG ATC ATC CAA G 6940

Trp Lys His Gln Asn Lys Asp Gln Asp Ile Val Lys Lys Ile Ile Gln

240

245

250

GTATGATAAT CTAAAATAAA AAGATCAATC AGAAATCAAA GACACCTATT TATCATAAAC 7000

CAGGAACAAG ACTGCATGTA TGTTTAGTTG TGTGGATCTT GTTCCCTGT TGAATCATT 7060

GTTGGACTGA AAAAGTTTCC ACCTGATAAT GTAGATGTGA TTCCACAAAC AGTTATACAA 7120

GGTTTTGTTT TCACCCCTGC TCCCAGTTT CCTTGTAAG TATGTTGAAC ACTCTAAGAG 7180

AAGAGAAATG CATTGAAGG CAGGGCTGTA TCTCAGGGAG TCGCTTCCAG ATCCCTTAAC 7240

GCTTCTGTAA GCAGCCCTC TAGACCACCA AGGAGAAGCT CTATAACCAC TTTGTATCTT 7300

ACATTGCACC TCTACCAAGA AGCTCTGTTG TATTTACTTG GTAATTCTCT CCAGGTAGGC 7360

TTTTCGTAGC TTACAAATAT GTTCTTATTA ATCCTCATGA TATGGCCTGC ATTAATAA 7420

TTTTAATGCC ATATGTTATG AGAATTAATG AGATAAAATC TGAAGAGTGT TTGAGCCTCT 7480

TGTAGGAAAA ACCTAGTTAC AGCAAAATGT TCTCACATCT TATAAGTTTA TATAAAGATT 7540

CTCCITTAGA AATGGTGTGA GAGAGAAACA GAGAGAGATA GGGAGAGAAG TGTGAAAGAA 7600

TCTGAAGAAA AGGAGTTTCA TCCAGTGTGG ACTGTAAGCT TTACGACACA TGATGGAAAG 7660

AGTTCTGACT TCAGTAAGCA TTGGGAGGAC ATGCTAGAAG AAAAAGGAAG AAGAGTTTCC 7720

ATAATGCAGA CAGGGTCAGT GAGAAATTCA TTCAGGTCCT CACCAGTAGT TAAATGACTG 7780

TATAGTCTTG CACTACCTTA AAAAATTCA AGTATCTGAA ACCGGGGCAA CAGATTTTAG 7840

GAGACCAACG TCTTTGAGAG CTGATTGCTT TTGCTTATCC AAAGAGTAAA CTTTATGTT 7900

TTGAGCAAAC CAAAAGTATT CTTTGAACGT ATAATTAGCC CTGAAGCCGA AAGAAAAGAG 7960

AAAATCAGAG ACCGTTAGAA TTGAAGCAA CCAAATCCC TATTTTATAA ATGAGGACAT 8020

5 TTTAACCAG AAAGATGAAC CGATTTGGCT TAGGGCTCAC AGATACTAAG TGAATCATGT 8080
 CATTAATAGA AATGTTAGTT CCTCCCTCTT AGGTTTGTAC CCTAGCTTAT TACTGAAATA 8140
 TTCTCTAGGC TGTGTGTCTC CTTTAGTTCC TCGACCTCAT GTCTTTGAGT TTTCAGATAT 8200
 CCTCCTCATG GAGGTAGTCC TCTGGTGCTA TGTGTATTCT TTAAAGGCTA GTTACGGCAA 8260
 10 TTAACCTATC AACTAGCGCC TACTAATGAA ACTTTGTATT ACAAAGTAGC TAACCTGAAT 8320
 ACTTTCCTTT TTTTCTGAAA TGTTATGGTG GTAATTTCTC AAACCTTTTC TTAGAAAAC 8380
 GAGAGTCATG TGTCTTATTT TCTACTGTTA ATTTTCAAAA TTAGGAGCTT CTTCCTAAAGT 8440
 TTTGTTGGAT GCCAAAAATA TATAGCATAT TATCTTATTA TAACAAAAAA TATTATCTC 8500
 AGTTCTTAGA AATAAATGGT GTCACCTAAC TCCCTCTCAA AAGAAAAGCT TATCATTGAA 8560
 20 ATATAATTAT GAAATTCTGC AAGAACCTTT TGCCTCAGGC TTGTTTTATG ATGCCATTGG 8620
 ATGAATATAA ATGATGTGAA CACTTATCTG GGCTTTTGCT TTATGCAG AT ATT GAC 8676

Asp Ile Asp

25
 30 CTC TGT GAA AAC AGC GTG CAG CGG CAC ATT GGA CAT GCT AAC CTC ACC 8724
 Leu Cys Glu Asn Ser Val Gln Arg His Ile Gly His Ala Asn Leu Thr
 255 260 265 270

35 TTC GAG CAG CTT CGT AGC TTG ATG GAA AGC TTA CCG GGA AAG AAA GTG 8772
 Phe Glu Gln Leu Arg Ser Leu Met Glu Ser Leu Pro Gly Lys Lys Val
 40 275 280 285

45 GGA GCA GAA GAC ATT GAA AAA ACA ATA AAG GCA TGC AAA CCC AGT GAC 8820
 Gly Ala Glu Asp Ile Glu Lys Thr Ile Lys Ala Cys Lys Pro Ser Asp
 290 295 300

50 CAG ATC CTG AAG CTG CTC AGT TTG TGG CGA ATA AAA AAT GCC GAC CAA 8868

55

Gln Ile Leu Lys Leu Leu Ser Leu Trp Arg Ile Lys Asn Gly Asp Gln

5

305

310

315

GAC ACC TTG AAG GGC CTA ATG CAC GCA CTA AAG CAC TCA AAG ACG TAC 8916

10

Asp Thr Leu Lys Gly Leu Met His Ala Leu Lys His Ser Lys Thr Tyr

320

325

330

15

CAC TTT CCC AAA ACT GTC ACT CAG AGT CTA AAG AAG ACC ATC AGG TTC 8964

His Phe Pro Lys Thr Val Thr Gln Ser Leu Lys Lys Thr Ile Arg Phe

20

335

340

345

350

25

CTT CAC AGC TTC ACA ATG TAC AAA TTG TAT CAG AAG TTA TTT TTA GAA 9012

Leu His Ser Phe Thr Met Tyr Lys Leu Tyr Gln Lys Leu Phe Leu Glu

355

360

365

30

ATG ATA GGT AAC CAG GTC CAA TCA GTA AAA ATA AGC TGC TTA 9054

Met Ile Gly Asn Gln Val Gln Ser Val Lys Ile Ser Cys Leu

35

370

375

380

40

TAACTGGAAA TGGCCATTGA GCTGTTTCCT CACAATTGGC GAGATCCCAT GGATGACTAA 9114

ACTGTTTCTC AGGCACTTGA GGCTTTCAGT GATATCTTTC TCATTACCAG TGAATAATT 9174

45

TGCCACAGGG TACTAAAAGA AACTATGATG TGGAGAAAGG ACTAACATCT CCTCCAATAA 9234

ACCCCAAATG GTTAATCCAA CTGTCAGATC TGGATCGTTA TCTACTGACT ATATTTTCCC 9294

TTATTACTGC TTGCAGTAAT TCAACTGGAA ATTAAAAAAA AAAAATACTAGA CTCCACTGGG 9354

50

CCTTACTAAA TATGGAATG TCTAACTTAA ATAGCTTTGG GATTCCAGCT ATGCTAGAGG 9414

CTTTTATTAG AAAGCCATAT TTTTCTCTGT AAAAGTTACT AATATATCTG TAACACTATT 9474

55

ACAGTATTGC TATTATATT CATTACAGATA TAAGATTTGG ACATATTATC ATCCTATAAA 9534
 5 GAAACGGTAT GACTTAATTT TAGAAAGAAA ATTATATTCT GTTTATTATG ACAAATGAAA 9594
 GAGAAAATAT ATATTTTAA TGGAAACTTT GTAGCATTTT TCTAATAGGT ACTGCCATAT 9654
 TTTTCTGTGT GGAGTATTTT TATAATTTTA TCTGTATAAG CTGTAATATC ATTTTATAGA 9714
 10 AAATGCATTA TTTAGTCAAT TGTTTAATGT TCGAAAACAT ATGAAATATA AATTATCTGA 9774
 ATATTAGATG CTCTGAGAAA TTGAATGTAC CTTATTTAAA AGATTTTATG GTTTTATAAC 9834
 TATATAAATG ACATTATTAA AGTTTTCAAA TTATTTTITA TTGCTTTCTC TGTGCTTTT 9894
 15 ATTT 9898

Sequence number: 3

20 Length of sequence: 401

Sequence Type: amino acid

Strandedness: single stranded

25 Topology: linear

Molecular type: protein

30

Sequence:

Met Asn Asn Leu Leu Cys Cys Ala Leu Val Phe Leu Asp Ile Ser
 35 -20 -15 -10
 Ile Lys Trp Thr Thr Gln Glu Thr Phe Pro Pro Lys Tyr Leu His
 40 -5 1 5
 Tyr Asp Glu Glu Thr Ser His Gln Leu Leu Cys Asp Lys Cys Pro
 10 15 20
 45 Pro Gly Thr Tyr Leu Lys Gln His Cys Thr Ala Lys Trp Lys Thr
 25 30 35
 50 Val Cys Ala Pro Cys Pro Asp His Tyr Tyr Thr Asp Ser Trp His
 40 45 50

55

	Ile Gln Asp Ile Asp Leu Cys Glu Asn Ser Val Gln Arg His Ile	
5	250	255 260
	Gly His Ala Asn Leu Thr Phe Glu Gln Leu Arg Ser Leu Met Glu	
10	265	270 275
	Ser Leu Pro Gly Lys Lys Val Gly Ala Glu Asp Ile Glu Lys Thr	
	280	285 290
15	Ile Lys Ala Cys Lys Pro Ser Asp Gln Ile Leu Lys Leu Leu Ser	
	295	300 305
20	Leu Trp Arg Ile Lys Asn Gly Asp Gln Asp Thr Leu Lys Gly Leu	
	310	315 320
25	Met His Ala Leu Lys His Ser Lys Thr Tyr His Phe Pro Lys Thr	
	325	330 335
30	Val Thr Gln Ser Leu Lys Lys Thr Ile Arg Phe Leu His Ser Phe	
	340	345 350
	Thr Met Tyr Lys Leu Tyr Gln Lys Leu Phe Leu Glu Met Ile Gly	
35	355	360 365
	Asn Gln Val Gln Ser Val Lys Ile Ser Cys Leu	
40	370	375 380

Sequence number: 4

45 Length of sequence: 1206

Sequence Type: nucleic acid

Strandedness: single stranded

50 Topology: linear

Molecular type: cDNA

55

Sequence:

5 ATGAACAAC TGTGTGCTG CGCGCTCGTG TTTCTGGACA TCTCCATTAA GTGGACCACC 60
 CAGGAAACGT TTCCTCCAAA GTACCTTCAT TATGACGAAG AACCTCTCA TCAGCTGTTG 120
 TGTGACAAAT GTCCTCCTGG TACCTACCTA AAACAACACT GTACAGCAAA GTGGAAGACC 180
 10 GTGTGCGCCC CTTGCCCTGA CCACTACTAC ACAGACAGCT GGCACACCAG TGACGACTGT 240
 CTATACTGCA GCCCGTGTG CAAGGAGCTG CAGTACGTC AGCAGGAGTG CAATCGCACC 300
 CACAACCGCG TGTGCGAATG CAAGGAAGGG CGCTACCTTG AGATAGAGTT CTGCTTGAAA 360
 15 CATAGGAGCT GCCCTCCTGG ATTTGGAGTG GTGCAAGCTG GAACCCGAGA GCGAAATACA 420
 GTTTGCAAAA GATGTCCAGA TGGTTCTTC TCAAATGAGA CGTCATCTAA AGCACCTGT 480
 20 AGAAAACACA CAAATTGCAG TGTCTTTGGT CTCCTGCTAA CTCAGAAAGG AAATGCAACA 540
 CACGACAACA TATGTCCGG AAACAGTGAA TCAACTCAA AATGTGGAAT AGATGTTACC 600
 25 CTGTGTGAGG AGGCATTCTT CAGGTTTGCT GTTCTTACAA AGTTTACGCC TAACTGGCTT 660
 AGTGTCTTGG TAGACAATTT GCCTGGCACC AAAGTAAACG CAGAGAGTGT AGAGAGGATA 720
 AAACGGCAAC ACAGCTCACA AGAACAGACT TTCCAGCTGC TGAAGTTATG GAAACATCAA 780
 30 AACAAAGACC AAGATATAGT CAAGAAGATC ATCCAAGATA TTGACCTCTG TCAAAACAGC 840
 GTGCAGCGGC ACATTGGACA TGCTAACCTC ACCTTCGAGC AGCTTCGTAG CTTGATGGAA 900
 35 AGCTTACCGG GAAAGAAAGT GCGAGCAGAA GACATTGAAA AAACAATAAA GGCATGCAAA 960
 CCCAGTGACC AGATCCTGAA GCTGCTCAGT TTGTGGCGAA TAAAAAATGG CGACCAAGAC 1020
 ACCTTGAAGG GCCTAATGCA CGCACTAAAG CACTCAAAGA CGTACCACTT TCCCAAAACT 1080
 40 GTCACTCAGA GTCTAAAGAA GACCATCAGG TTCCTTCACA GCTTCACAAT GTACAAATTG 1140
 TATCAGAAGT TATTTTLAGA AATGATAGGT AACCAGGTCC AATCAGTAAA AATAAGCTGC 1200
 45 TTATAA 1206

50

55

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: SNOW BRAND MILK PRODUCTS CO., LTD.
 (B) STREET: 1-1, NAEBOCHO 6-CHOME
 (C) CITY: HIGASHI-KU, SAPPORO-SHI
 (D) STATE: HOKKAIDO
 (E) COUNTRY: JP
 (F) POSTAL CODE (ZIP): NONE

(ii) TITLE OF INVENTION: NOVEL DNA AND PROCESS FOR PREPARING PROTEIN USING THE DNA

(iii) NUMBER OF SEQUENCES: 4

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: EP 97935810.8

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: JP 235928/96
 (B) FILING DATE: 19-AUG-1996

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1316 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: genomic DNA (human OCIF genomic DNA-1)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

CTGGAGACAT ATAACITGAA CACTTGGCCC TGATGGGGAA GCAGCTCTGC AGGGACTTTT 60
TCAGCCATCT GTAACAATT TCAGTGGCAA CCGCGAACT GTAATCCATG AATGGGACCA 120
CACTTTACAA GTCATCAAGT CTAACITCTA GACCAGGGAA TTAATGGGGG AGACAGCGAA 180
CCCTAGAGCA AAGTGCCAAA CTTCTGTCTG TAGCTTGAGG CTAGTGGAAG GACCTCGAGG 240
AGGCTACTCC AGAAGTTCAG CGCGTAGGAA GCTCCGATAC CAATAGCCCT TTGATGATGG 300
TGGGGTTGGT GAAGGGAACA GTGCTCCGCA AGGTTATCCC TGCCCCAGGC AGTCCAATTT 360
TCACTCTGCA GATTCTCTCT GGCTCTAACT ACCCCAGATA ACAAGGAGTG AATGCAGAAT 420
AGCACGGGCT TTAGGGCCAA TCAGACATTA GTTAGAAAAA TTCCTACTAC ATGGTTTATG 480
TAAACTTGAA GATGAATGAT TGCGAACTCC CCGAAAAGGG CTCAGACAAT GCCATGCATA 540
AAGAGGGGCC CTGTAATTTG AGGTTTCAGA ACCCGAAGTG AAGGGGTGAG GCAGCCGGGT 600
ACGGCGGAAA CTCACAGCTT TCGCCAGCG AGAGGACAAA GGTCTGGGAC AACTCCAAC 660
TGCGTCCGGA TCTTGGCTGG ATCGGACTCT CAGGGTGGAG GAGACACAAG CACAGCAGCT 720
GCCCAGCGTG TGCCAGCCCC TCCCACCGCT GGTCCCGGCT GCCAGGAGGC TGGCCGCTGG 780
CGGGAAGGGG CCGGGAACCC TCAGAGCCCC GCGGAGACAG CAGCCGCCCT GTTCTCTAGC 840
CCGGTGGCTT TTTTTCGCC TGCTCTCCCA GGGGACAGAC ACCACCGCCC CACCCCTCAC 900
GCCCCACCTC CTTGGGGGAT CCTTTCCGCG CCAGCCCTGA AAGCGTTAAT CCTGGAGCTT 960
TCTGCACACC CCCCAGCCGC TCCCAGCCAA GCTTCTTAAA AAAGAAAGGT GCAAAGTTTG 1020
GTCCAGGATA GAAAAATGAC TGATCAAAGG CAGGCGATAC TTCTGTGTC CGGGACGCTA 1080
TATATAACGT GATGACGCA CGGGCTGCGG AGACGCACCG GAGCGCTCGC CCAGCCGCCG 1140
CCTCCAAGCC CTGAGGTTT CCGGGGACCA CA ATG AAC AAG TTG CTG TGC TGC 1193
Met Asn Lys Leu Leu Cys Cys
-20 -15

```

GCG CTC GTG GTAAGTCCT GGGCCAGCCG ACGGGTCCCC GCGCCTGGG 1242

Ala Leu Val

GAGGCTGCTG CCACCTGGTC TCCCAACCTC CCAGCGGACC GCGGGGAGA AGGCTCCACT 1302
CGCTCCCTCC CAGG 1316

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9898 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: genomic DNA (human OCIF genomic DNA-2)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GCTTACTTTG TGCCAAATCT CATTAGGCTT AAGGTAATAC AGGACTTTGA GTCAAATGAT 60
ACTGTTGCAC ATAAGAACA ACCTATTTTC ATGCTAAGAT GATGCCACTG TGTTCTTTTC 120
TCCTTCTAG TTT CTG GAC ATC TCC ATT AAG TGG ACC ACC CAG GAA ACG TTT 171
Phe Leu Asp Ile Ser Ile Lys Trp Thr Thr Gln Glu Thr Phe
-10 -5 1

OCT CCA AAG TAC CTT CAT TAT GAC GAA GAA ACC TCT CAT CAG CTG TTG 219
Pro Pro Lys Tyr Leu His Tyr Asp Glu Glu Thr Ser His Gln Leu Leu
5 10 15

TGT GAC AAA TGT CCT CCT GGT ACC TAC CTA AAA CAA CAC TGT ACA GCA 267
Cys Asp Lys Cys Pro Pro Gly Thr Tyr Leu Lys Gln His Cys Thr Ala
20 25 30 35

AAG TGG AAG ACC GTG TGC GCC CCT TGC CCT GAC CAC TAC TAC ACA GAC 315
Lys Trp Lys Thr Val Cys Ala Pro Cys Pro Asp His Tyr Tyr Thr Asp
40 45 50

AGC TGG CAC ACC AGT GAC GAG TGT CTA TAC TGC AGC CCC GTG TGC AAG 363
Ser Trp His Thr Ser Asp Glu Cys Leu Tyr Cys Ser Pro Val Cys Lys
55 60 65

GAG CTG CAG TAC GTC AAG CAG GAG TGC AAT CGC ACC CAC AAC CGC GTG 411
Glu Leu Gln Tyr Val Lys Gln Glu Cys Asn Arg Thr His Asn Arg Val
70 75 80

TGC GAA TGC AAG GAA GGG CGC TAC CTT GAG ATA GAG TTC TGC TTG AAA 459
Cys Glu Cys Lys Glu Gly Arg Tyr Leu Glu Ile Glu Phe Cys Leu Lys
85 90 95

CAT AGG AGC TGC CCT CCT GGA TTT GGA GTG GTG CAA GCT G GTACGTGTCA 509
His Arg Ser Cys Pro Pro Gly Phe Gly Val Val Gln Ala
100 105 110

ATGTGCAGCA AAATTAATTA GGATCATGCA AAGTCAGATA GTTGTGACAG TTAGGAGAA 569
CACTTTTGTT CTGATGACAT TATAGGATAG CAAATTGCAA AGGTAATGAA ACCTGCCAGG 629
TAGTACTAT GTGCTGGAG TGCTTCCAAA GGACCATTCG TCAGAGGAAT ACTTTGCCAC 689
TACAGGGCAA TTTAATGACA AATCTCAAA GCAGCAAATT ATTCTCTCAT GAGATGCATG 749
ATGGTTTTTT TTTTTTTTTT TAAAGAAACA AACTCAAGTT GCACTATTGA TAGTTGATCT 809
ATACCTCTAT ATTCACTTC AGCATGGACA CCTTCAAAC GCAGCACTTT TTGACAAACA 869
TCAGAAATGT TAATTTATAC CAAGAGAGTA ATTATGCTCA TATTAATGAG ACTCTGGAGT 929
GCTAACATA AGCAGTTATA ATTAATTATG TAAAAATGA GAATGGTGAG GGAATTGCA 989
TTTCATTATT AAAACAAGG CTAGTTCCTC CTTAGCATG GGAGCTGAGT GTTTGGGAGG 1049
GTAAGGACTA TAGCAGAATC TCTTCAATGA GCTTATCTT TATCTTAGAC AAAACAGATT 1109
GTCAAGCCAA GAGCAAGCAC TTGCCTATAA ACCAAGTGCT TTCTCTTTTG CATTTTGAAC 1169
AGCATTGGTC AGGGCTCATG TGTATTGAAT CTTTAAACC AGTAACCCAC GTTTTTTTTC 1229
TGCCACATTT GCGAAGCTTC AGTCAGCCT ATAACCTTTT ATAGCTTGAG AAAATTAAGA 1289
GTATCCACTT ACTTAGATGG AAGAAGTAAT CAGTATAGAT TCTGATGACT CAGTTTGAAG 1349

	CAGTGTCTCT	CAACTGAAGC	CCTGCTGATA	TTTAAAGAAA	TATCTGGATT	CCTAGGCTGG	1409
	ACTCCTTTTT	GTGGGCAGCT	GTCCCTGCGCA	TTGTAGAATT	TTGGCAGCAC	CCCTGGACTC	1469
	TAGCCACTAG	ATACCAATAG	CAGTCCTTCC	CCCATGTGAC	AGCCAAAAAT	GTCTTCAGAC	1529
5	ACTGTCAAAT	GTGCCCAGGT	GGCAAAATCA	CTCCTGGTTG	AGAACAGGGT	CATCAATGCT	1589
	AAGTATCTGT	AACTATTTTA	ACTCTCAAAA	CTTGTGATAT	ACAAAAGCTA	AATTATTAGA	1649
	CGACCAATAC	TTTAGGTTTA	AAGGCATACA	AATGAAACAT	TCAAAAATCA	AAATCTATTG	1709
	TGTTTCTCAA	ATAGTGAATC	TTATAAAATT	AATCACAGAA	GATGCAAATT	GCATCAGAGT	1769
	CCCTTAAAAAT	TCTCTTTCGT	ATGAGTATTT	GAGGGAGGAA	TTGGTGATAG	TTCTTACTTT	1829
	CTATTGGATG	GTACTTTGAG	ACTCAAAAGC	TAAGCTAAGT	TGTGTGTGTG	TCAGGGTGCG	1889
10	GGGTGTGGAA	TCCCATCAGA	TAAAAAGCAA	TCCATGTAAT	TCATTCAGTA	AGTTGTATAT	1949
	GTAGAAAAAT	GAAAAGTGGG	CTATGCAGCT	TGGAACTAG	AGAATTTTGA	AAAATAATGG	2009
	AAATCACAAG	GATCTTTCTT	AAATAAGTAA	GAAAATCTGT	TTGTAGAATG	AAGCAAGCAG	2069
	GCAGCCAGAA	GACTCAGAAC	AAAAGTACAC	ATTTTACTCT	GTGTACACTG	GCAGCACAGT	2129
	GGGATTTATT	TACCTCTCCC	TCCCTAAAAA	CCCACACAGC	GGTTCCTCTT	GGGAAATAAG	2189
	AGGTTTCCAG	CCCAAAGAGA	AGGAAAGACT	ATGTGGTGTG	ACTCTAAAAA	GTATTTAATA	2249
	ACCGTTTTGT	TGTTGCTGTT	GCTGTTTTGA	AATCAGATTG	TCTCCTCTCC	ATATTTTATT	2309
15	TACTTCATTC	TGTTAATTCC	TGTGGAATTA	CTTAGAGCAA	GCATGGTGAA	TTCTCAACTG	2369
	TAAAGCCAAA	TTTCTCCATC	ATTATAATTT	CACATTTTGC	CTGGCAGGTT	ATAATTTTTA	2429
	TATTTCCACT	GATAGTAATA	AGGTAAATAT	ATTACTTAGA	TGGATAGATC	TTTTTCATAA	2489
	AAAGTACCAT	CAGTTATAGA	GGGAAGTCAT	GTTTCATGTT	AGGAAGGTCA	TTAGATAAAG	2549
	CTTCTGAATA	TATTTATGAA	CATTAGTTCT	GTCATTCTTA	GATTCTTTT	GTTAAATAAC	2609
	TTTAAAAGCT	AACCTACCTA	AAAGAAATAT	CTGACACATA	TGAACCTCTC	ATTAGGATGC	2669
20	AGGAGAAGAC	CCAAGCCACA	GATATGTATC	TGAAGAATGA	ACAAGATTCT	TAGGCCCGGC	2729
	ACGGTGGCTC	ACATCTGTAA	TCTCAAGAGT	TTGAGAGGTC	AAGGCCGGCA	GATCACCTGA	2789
	GGTCAGGAGT	TCAAGACCAG	CCTGGCCAAC	ATGATGAAAC	CCTGCCTCTA	CTAAAAATAC	2849
	AAAAATTAGC	AGGCAATGGT	GGTGCATGCC	TGCAACCCCTA	GCTACTCAGG	AGGCTGAGAC	2909
	AGGAGAACTC	CTTGAACCCCT	CGAGGCGGAG	GTTGTGGTGA	GCTGAGATCC	CTCTACTGCA	2969
	CTCCAGCCTG	GGTGACAGAG	ATGAGACTCC	GTCCCTGCCG	CCGCCCCCGC	CTTCCCCCCC	3029
25	AAAAAGATTC	TTCTTCATGC	AGAACATACG	GCAGTCAACA	AAGGGAGACC	TGGGTCCAGG	3089
	TGTCCAAGTC	ACTTATTTTCG	AGTAAATTAG	CAATGAAAGA	ATGCCATGGA	ATCCCTGCCC	3149
	AAATACCTCT	GCTTATGATA	TTGTAGAATT	TGATATAGAG	TTGTATCCCA	TTTAAGGAGT	3209
	AGGATGTAGT	AGGAAAGTAC	TAAAAACAAA	CACACAAACA	GAAAACCCCTC	TTTGCTTTGT	3269
	AAGGTGGTTC	CTAAGATAAT	GTCAAGTGCAA	TGCTGGAAAT	AATATTTAAT	ATGTGAAGGT	3329
	TTTAGGCTGT	GTTTTCCTCT	CCTGTCTCTT	TTTTCTGCCA	GCCCTTTGTC	ATTTTTCAG	3389
30	GTCAATGAAT	CATGTAGAAA	GAGACAGGAG	ATGAAACTAG	AACCAGTCCA	TTTTGCCCTC	3449
	TTTTTTATTT	TCTGGTTTTG	GTAAAAAGATA	CAATGAGGTA	GGAGGTTGAG	ATTTATAAAT	3509
	GAAGTTTAAT	AAGTTTCTGT	AGCTTTGATT	TTTCTCTTTC	ATATTFTGTA	TCTTGACATA	3569
	GCCAGAAATG	GCCTGTAAAA	TCTACATATG	GATATTGAAG	TCTAAATCTG	TTCAACTAGC	3629
	TTACACTAGA	TGGGATATT	TTCATATTCA	GATACACTGG	AATGTATGAT	TAGCCATGTC	3689
	GTAATATAGT	CAAGTGTTTG	AAGGTATTTA	TTTTTAATAG	CGTCTTTAGT	TGTGGACTGG	3749
	TTCAAGTTTT	TCTGCCAATG	ATTTCTTCAA	ATTTATCAAA	TATTTTTTCA	TCATGAAGTA	3809
35	AAATGCCCTT	GCAGTCACCC	TTCCGTGAAGT	TTGAACGACT	CTGCTGTTTT	AAACAGTTTA	3869
	AGCAAAATGGT	ATATCATCTT	CCGTTTACTA	TGTAGCTTAA	CTGCAGGCTT	ACGCTTTTGA	3929
	GTCAGCGGCC	AACTTTATTG	CCACCTTCAA	AAGTTTATTA	TAATGTTGTA	AATTTTTTACT	3989
	TCTCAAGGTT	AGCATACTTA	GGAGTTGCTT	CACAATTAGG	ATTCAGGAAA	GAAAGAACTT	4049
	CAGTAGGAAC	TGATTGGAAT	TTAATGATGC	AGCATTCAAT	GGGTACTAAT	TTCAAGAAT	4109
	GATATTACAG	CAGACACACA	GCAGTTATCT	TGATTTTCTA	GGAATAATTG	TATGAAGAAT	4169
40	ATGGCTGACA	ACACGGCCTT	ACTGCCACTC	AGCGGAGGCT	GGACTAATGA	ACACCCCTACC	4229
	CTTCTTTCTT	TTCTCTCTAC	ATTTCTATGAG	CGTTTTGTAG	GTAACGAGAA	AATTGACTTG	4289
	CATTTGCATT	ACAAGGAGGA	GAAACTGGCA	AAGGGGATGA	TGGTGGAAGT	TTTGTCTCTG	4349
	CTAATGAAGT	GAAAAATGAA	AATGCTAGAG	TTTTGTGCAA	CATAATAGTA	GCAGTAAAAA	4409
	CCAAGTGAAA	AGTCTTTCCA	AAACTGTGTT	AAGAGGGCAT	CTGCTGGGAA	ACGATTTGAG	4469
	GAGAAGGTAC	TAAATTGCTT	GGTATTTTCC	GTAG GA ACC	CCA GAG CGA	AAT ACA	4523
45				Gly Thr Pro Glu Arg Asn Thr			
				115			
	GTT TGC AAA AGA TGT CCA GAT GGG TTC TTC TCA AAT GAG ACG TCA TCT	4571					
	Val Cys Lys Arg Cys Pro Asp Gly Phe Phe Ser Asn Glu Thr Ser Ser						
	120 125 130 135						
50	AAA GCA CCC TGT AGA AAA CAC ACA AAT TGC AGT GTC TTT GGT CTC CTG	4619					
	Lys Ala Pro Cys Arg Lys His Thr Asn Cys Ser Val Phe Gly Leu Leu						
	140 145 150						
	CTA ACT CAG AAA GGA AAT GCA ACA CAC GAC AAC ATA TGT TCC GGA AAC	4667					
55							

EP 0 874 045 A1

Leu Thr Gln Lys Gly Asn Ala Thr His Asp Asn Ile Cys Ser Gly Asn
155 160 165

AGT GAA TCA ACT CAA AAA TGT GGA ATA G GTAATTACAT TCCAAAATAC 4715
5 Ser Glu Ser Thr Gln Lys Cys Gly Ile
170 175

GTCTTTGTAC GATTTTGTAG TATCATCTCT CTCTCTGAGT TGAACACAAG GCCTCCAGCC 4775
ACATTTCTTGG TCAAACTTAC ATTTTCCCTT TCTTGAATCT TAACCAGCTA AGGCTACTCT 4835
10 CGATGCATTA CTGCTAAAGC TACCACTCAG AATCTCTCAA AAATCTCATCT TCTCAGAGAT 4895
AACACCTCAA AGCTTGATTT TCTCTCCTTT CACACTGAAA TCAAAATCTTG CCCATAGGCA 4955
AAGGGCAGTG TCAAGTTTGC CACTGAGATG AAATTAGGAG AGTCCAAACT GTAGAATTCA 5015
CGTTGTGTGT TATTACTTTC ACGAATGTCT GTATTATTAA CTAAAGTATA TATTGGCAAC 5075
TAAGAAGCAA AGTGATATAA ACATGATGAC AAATTAGGCC AGGCATGGTG GCTTACTCCT 5135
ATAATCCCAA CATTTTGGGG GGCCAAAGTA GGCAGATCAC TTGAGGTCAG GATTTCAAGA 5195
CCAGCCTGAC CAACATGGTG AAACCTTGTC TCTACTAAAA ATACAAAAAT TAGCTGGGCA 5255
15 TGSTAGCAGG CACTTCTAGT ACCAGCTACT CAGGGCTGAG GCAGGAGAAT CGCTTGAACC 5315
CAGGAGATGG AGGTTCAGT GAGCTGAGAT TGTACCACTG CACTCCAGTC TGGGCAACAG 5375
AGCAAGATTG CATCACACAC ACACACACAC ACACATTAGA AATGTGTACT 5435
TGGCTTTGTT ACCTATGGTA TTAGTGATC TATTGCATGG AACTTCCAAG CTACTCTGGT 5495
TGTGTTAAGC TCTTCATTGG GTACAGGTCA CTAGTATTAA GTTCAGGTTA TTCGGATGCA 5555
TTCCACGGTA GTGATGACAA TTTCATCAGG TAGTGTGTGT GTTCACCTTG TCACTCCAC 5615
20 CACTAGACTA ATCTCAGACC TTCACTCAA GACACATTAC ACTAAAGATG ATTTGCTTTT 5675
TTGTGTTTAA TCAAGCAATG GTATAAACCA GCCTGACTCT CCCCAAACAG TTTTTCGTAC 5735
TACAAAGAAG TTTATGAAGC AGAGAAATGT GAATTGATAT ATATATGAGA TTCTAACCCA 5795
GTTCCAGCAT TGTTCATTG TGTAAATTGAA ATCATAGACA AGCCATTTTA GCCTTTGCTT 5855
TCTTATCTAA AAAAAAAAAA AAAAAAATGA AGGAAGGGGT ATTAAGAGGA GTGATCAAAT 5915
TTTAACATTC TCTTTAATTA ATTCATTTTT AATTTTACTT TTTTTCATTT ATTGTGCACT 5975
25 TACTATGTGG TACTGTGCTA TAGAGGCTTT AACATTTATA AAAACACTGT GAAAGTTGCT 6035
TCAGATGAAT ATAGGTAGTA GAACGGCAGA ACTAGTATTC AAAGCCAGGT CTGATGAATC 6095
CAAAAACAAA CACCCATTAC TCCCATTTC TGGGACATAC TTACTCTACC CAGATGCTCT 6155
GGGCTTTGTA ATGCCATGT AAATAACATA GTTTTATGTT TGGTTATTTT CCTATGTAAT 6215
GTCTACTTAT ATATCTGTAT CTATCTCTTG CTTTGTTCCT AAAGGTAAAC TATGTGTCTA 6275
AATGTGGGCA AAAAAATACA CACTATTCCA AATTACTGTT CAAATTCCTT TAAGTCAGTG 6335
30 ATAATTATTT GTTTTGACAT TAATCATGAA GTTCCCTGTG GGTACTAGGT AAACCTTTAA 6395
TAGAATGTTA ATGTTTGTAT TCATTATAAG AATTTTGGC TGTACTTAT TTACAACAAT 6455
ATTTCACTCT AATTAGACAT TTACTAAACT TTCTCTTGAA AACATGCCC AAAAAAGAAC 6515
ATTAGAAGAC ACGTAAGCTC AGTTGGTCTC TGCCACTAAG ACCAGCCAAC AGAAGCTTGA 6575
TTTATTTCAA ACTTTGCATT TTAGCATATT TTATCTTGGA AAATTCATT GTGTTGGTTT 6635
TTTGTTTTGG TTTGTATTGA ATAGACTCTC AGAAATCCAA TTGTTGAGTA AATCTTCTGG 6695
GTTTCTAAC CTTTCTTTAG AT GTT ACC CTG TGT GAG GAG GCA TTC TTC AGG 6747
35 Asp Val Thr Leu Cys Glu Glu Ala Phe Phe Arg
180 185

TTT GCT GTT CCT ACA AAG TTT ACG CCT AAC TGG CTT AGT GTC TTG GTA 6795
Phe Ala Val Pro Thr Lys Phe Thr Pro Asn Trp Leu Ser Val Leu Val
190 195 200

GAC AAT TTG CCT GGC ACC AAA GTA AAC GCA GAG AGT GTA GAG AGG ATA 6843
40 Asp Asn Leu Pro Gly Thr Lys Val Asn Ala Glu Ser Val Glu Arg Ile
205 210 215

AAA CGG CAA CAC AGC TCA CAA GAA CAG ACT TTC CAG CTG CTG AAG TTA 6891
45 Lys Arg Gln His Ser Ser Gln Glu Gln Thr Phe Gln Leu Leu Lys Leu
220 225 230 235

TGG AAA CAT CAA AAC AAA GAC CAA GAT ATA GTC AAG AAG ATC ATC CAA G 6940
Trp Lys His Gln Asn Lys Asp Gln Asp Ile Val Lys Lys Ile Ile Gln
240 245 250

GTATGATAAT CTAAATAAAA AAGATCAATC AGAAATCAAA GACACCTATT TATCATAAAC 7000
50 CAGGAACAAG ACTGCATGTA TGTTTAGTTG TGTGGATCTT GTTCCCTGT TGGAAATCATT 7060
GTTGGACTGA AAAAGTTTCC ACCTGATAAT GTAGATGTGA TTCCACAAC AGTTATACAA 7120
GGTTTGTTC TCACCCCTGC TCCCAGTTT CCTTGTAAG TATGTTGAAC ACTCTAAGAG 7180
AAGAGAAATG CATTTGAAGG CAGGGCTGTA TCTCAGGGAG TCGCTTCCAG ATCCCTTAAC 7240

EP 0 874 045 A1

5 GCTTCTGTAA GCAGCCCCCTC TAGACCACCA AGGAGAAGCT CTATAACCAC TTTGTATCTT 7300
 ACATTGCACC TCTACCAAGA AGCTCTGTGT TATTACTTGT GTAATTCTCT CCAGGTAGGC 7360
 TTTTCGTAGC TTACAAATAT GTTCTTATTA ATCCTCATGA TATGGCCTGC ATTAAAAATTA 7420
 TTTAATGGC ATATGTTATG AGAATTAATG AGATAAAATC TGAAAAGTGT TTGAGCCTCT 7480
 TGTAGGAAAA AGCTAGTTAC AGCAAAATGT TCTCACATCT TATAAGTTTA TATAAGATT 7540
 CTCCTTTAGA AATGGTGTGA GAGAGAAACA GAGAGAGATA GGGAGAGAAG TGTGAAAGAA 7600
 TCTGAAGAAA AGGAGTTTCA TCCAGTGTGG ACTGTAAGCT TTACGACACA TGATGGAAAG 7660
 AGTTCTGACT TCAGTAAGCA TTGGGAGGAC ATGCTAGAAG AAAAAGGAAG AAGAGTTTCC 7720
 ATAATGCAGA CAGGGTCAGT GAGAAATTC AATCAGGTCT CACCAGTAGT TAAATGACTG 7780
 10 TATAGTCTTG CACTACCCTA AAAAATCTCA AGTATCTGAA ACCGGGGCAA CAGATTTTAG 7840
 GAGACCAACG TCTTTGAGAG CTGATTGCTT TTGCTTATGC AAAGAGTAAA CTTTATGTT 7900
 TTGAGCAAAC CAAAAGTATT CTTTGAACTG ATAATTAGCC CTGAGCCGA AAGAAAAGAG 7960
 AAAATCAGAG ACCGTTAGAA TTGGAAGCAA CCAAATTCCT TATTTTATAA ATGAGGACAT 8020
 TTAAACCCAG AAAGATGAAC CGATTGGCT TAGGGCTCAC AGATACTAAG TGACTCATGT 8080
 CATTAATAGA AATGTTAGTT CCTCCCTCTT AGGTTTGTAC CCTAGCTTAT TACTGAAATA 8140
 TTTCTTAGGC TGTGTGCTC CTTTAGTTC TCGACCTCAT GTCTTTGAGT TTTGAGATAT 8200
 15 CCTCCTCATG GAGGTAGTCC TCTGGTGCTA TGTGTATTCT TTAAGGGCTA GTTACGGCAA 8260
 TTAACCTATC AACTAGCGCC TACTAATGAA ACTTTGTATT ACAAAGTAGC TAACTTGAAT 8320
 ACTTTCTTTT TTTTCTGAAA TGTATGGTG GTAATTTCTC AAATTTTTC TTAGAAAAC 8380
 GAGAGTGATG TGTCTTATTT TCTACTGTTA ATTTTCAAAA TTAGGAGCTT CTTCCAAAGT 8440
 TTTGTTGGAT GCCAAAAATA TATAGCATAT TATCTTATTA TAACAAAAAA TATTATCTC 8500
 AGTTCTTAGA AATAAATGGT GTCACTTAAC TCCCTCTCAA AAGAAAAGGT TATCATTGAA 8560
 20 ATATAATTAT GAAATCTGC AAGAACCTTT TGCCTCACGC TTGTTTATG ATGGCATTGG 8620
 ATGAATATAA ATGATGTGAA CACTTATCTG GGCTTTTGT TATGTCAG AT ATT GAC 8676
 Asp Ile Asp
 CTC TGT GAA AAC AGC GTG CAG CGG CAC ATT GGA CAT GCT AAC CTC ACC 8724
 Leu Cys Glu Asn Ser Val Gln Arg His Ile Gly His Ala Asn Leu Thr
 25 255 260 265 270
 TTC GAG CAG CTT CGT AGC TTG ATG GAA AGC TTA CCG GGA AAG AAA GTG 8772
 Phe Glu Gln Leu Arg Ser Leu Met Glu Ser Leu Pro Gly Lys Lys Val
 275 280 285
 GGA GCA GAA GAC ATT GAA AAA ACA ATA AAG GCA TGC AAA CCC AGT GAC 8820
 Gly Ala Glu Asp Ile Glu Lys Thr Ile Lys Ala Cys Lys Pro Ser Asp
 290 295 300
 CAG ATC CTG AAG CTG CTC AGT TTG TGG CGA ATA AAA AAT GGC GAC CAA 8868
 Gln Ile Leu Lys Leu Leu Ser Leu Trp Arg Ile Lys Asn Gly Asp Gln
 305 310 315
 35 GAC ACC TTG AAG GGC CTA ATG CAC GCA CTA AAG CAC TCA AAG ACG TAC 8916
 Asp Thr Leu Lys Gly Leu Met His Ala Leu Lys His Ser Lys Thr Tyr
 320 325 330
 CAC TTT CCC AAA ACT GTC ACT CAG AGT CTA AAG AAG ACC ATC AGG TTC 8964
 His Phe Pro Lys Thr Val Thr Gln Ser Leu Lys Lys Thr Ile Arg Phe
 40 335 340 345 350
 CTT CAC AGC TTC ACA ATG TAC AAA TTG TAT CAG AAG TTA TTT TTA GAA 9012
 Leu His Ser Phe Thr Met Tyr Lys Leu Tyr Gln Lys Leu Phe Leu Glu
 355 360 365
 45 ATG ATA GGT AAC CAG GTC CAA TCA GTA AAA ATA AGC TGC TTA 9054
 Met Ile Gly Asn Gln Val Gln Ser Val Lys Ile Ser Cys Leu
 370 375 380
 50 TAACTGGAAA TGGCCATTGA GCTGTTTCCT CACAATTGGC GAGATCCCAT GGATGAGTAA 9114
 ACTGTTTCTC AGGCACCTGA GGCTTTTCAGT GATATCTTTC TCATTACCAG TGACTAATTT 9174
 TGCCACAGGG TACTAAAAGA AACTATGATG TGGAGAAAGG ACTAACATCT CCTCCAATAA 9234
 ACCCCAAATG GTTAATCCAA CTGTCAGATC TGGATCGTTA TCTACTGACT ATATTTTCCC 9294
 TTATTACTGC TTGCACTAAT TCAACTGGAA ATTAATAAAA AAAAACTAGA CTCCACTGGG 9354
 CCTTACTAAA TATGGGAATG TCTAACTTAA ATAGCTTTGG GATTCCAGCT ATGCTAGAGG 9414
 CTTTATATAG AAAGCCATAT TTTTTCCTGT AAAAGTTACT AATATATCTG TAACACTATT 9474
 55

ACAGTATTGC TATTTATATT CATTGAGATA TAAGATTGG ACATATTATC ATCCTATAAA 9534
 GAAACGGTAT GACTTAATTT TAGAAAGAAA ATTATATTCT GTTTATTATG ACAAATGAAA 9594
 GAGAAAATAT ATATTTTAA TGGAAAGTTT GTAGCATTMT TCTAATAGGT ACTGCCATAT 9654
 TTTTCGTGT GGAGTATTTT TATAATTTTA TCTGTATAAG CTGTAATATC ATTTTATAGA 9714
 AAATGCATTA TTAGTCAAT TGTTTAATGT TGGAAAACAT ATGAAATATA AATTATCTGA 9774
 ATATTAGATG CTCTGAGAAA TTGAATGTAC CTTATTTAAA AGATTTTATG GTTTTATAAC 9834
 TATATAAATG ACATTATTAA AGTTTTCAAA TTATTTTAA TTGCTTCTC TGTGCTTTT 9894
 ATTT

10

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 401 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

20

25

30

35

40

45

50

55

Met Asn Asn Leu Leu Cys Cys Ala Leu Val Phe Leu Asp Ile Ser
 -20 -15 -10
 Ile Lys Trp Thr Thr Gln Glu Thr Phe Pro Pro Lys Tyr Leu His
 -5 1 5
 Tyr Asp Glu Glu Thr Ser His Gln Leu Leu Cys Asp Lys Cys Pro
 10 15 20
 Pro Gly Thr Tyr Leu Lys Gln His Cys Thr Ala Lys Trp Lys Thr
 25 30 35
 Val Cys Ala Pro Cys Pro Asp His Tyr Tyr Thr Asp Ser Trp His
 40 45 50
 Thr Ser Asp Glu Cys Leu Tyr Cys Ser Pro Val Cys Lys Glu Leu
 55 60 65
 Gln Tyr Val Lys Gln Glu Cys Asn Arg Thr His Asn Arg Val Cys
 70 75 80
 Glu Cys Lys Glu Gly Arg Tyr Leu Glu Ile Glu Phe Cys Leu Lys
 85 90 95
 His Arg Ser Cys Pro Pro Gly Phe Gly Val Val Gln Ala Gly Thr
 100 105 110
 Pro Glu Arg Asn Thr Val Cys Lys Arg Cys Pro Asp Gly Phe Phe
 115 120 125
 Ser Asn Glu Thr Ser Ser Lys Ala Pro Cys Arg Lys His Thr Asn
 130 135 140
 Cys Ser Val Phe Gly Leu Leu Leu Thr Gln Lys Gly Asn Ala Thr
 145 150 155
 His Asp Asn Ile Cys Ser Gly Asn Ser Glu Ser Thr Gln Lys Cys
 160 165 170
 Gly Ile Asp Val Thr Leu Cys Glu Glu Ala Phe Phe Arg Phe Ala
 175 180 185
 Val Pro Thr Lys Phe Thr Pro Asn Trp Leu Ser Val Leu Val Asp
 190 195 200
 Asn Leu Pro Gly Thr Lys Val Asn Ala Glu Ser Val Glu Arg Ile
 205 210 215
 Lys Arg Gln His Ser Ser Gln Glu Gln Thr Phe Gln Leu Leu Lys
 220 225 230
 Leu Trp Lys His Gln Asn Lys Asp Gln Asp Ile Val Lys Lys Ile
 235 240 245
 Ile Gln Asp Ile Asp Leu Cys Glu Asn Ser Val Gln Arg His Ile
 250 255 260
 Gly His Ala Asn Leu Thr Phe Glu Gln Leu Arg Ser Leu Met Glu
 265 270 275
 Ser Leu Pro Gly Lys Lys Val Gly Ala Glu Asp Ile Glu Lys Thr
 280 285 290
 Ile Lys Ala Cys Lys Pro Ser Asp Gln Ile Leu Lys Leu Leu Ser
 295 300 305

Leu Trp Arg Ile Lys Asn Gly Asp Gln Asp Thr Leu Lys Gly Leu
 310 315 320
 Met His Ala Leu Lys His Ser Lys Thr Tyr His Phe Pro Lys Thr
 5 325 330 335
 Val Thr Gln Ser Leu Lys Lys Thr Ile Arg Phe Leu His Ser Phe
 340 345 350
 Thr Met Tyr Lys Leu Tyr Gln Lys Leu Phe Leu Glu Met Ile Gly
 355 360 365
 10 Asn Gln Val Gln Ser Val Lys Ile Ser Cys Leu
 370 375 380

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1206 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ATGAACAACT TGCTGTGCTG CGCGCTCGTG TTTCTGGACA TCTCCATTAA GTGGACCACC 60
 25 CAGGAAACGT TTCCTCCAAA GTACCTTCAT TATGACGAAG AAACCTCTCA TCAGCTGTTG 120
 TGTGACAAAT GTCCTCCTGG TACCTACCTA AAACAACACT GTACAGCAAA GTGGAAGACC 180
 GTGTGCGCCC CTTGCCCTGA CCACTACTAC ACAGACAGCT GGCACACCAG TGACGAGTGT 240
 CTATACTGCA GCCCCGTGTG CAAGGAGCTG CAGTACGTCA AGCAGGAGTG CAATCGCACC 300
 CACAACCGCG TGTGCGAATG CAAGGAAGGG CGCTACCTTG AGATAGAGTT CTGCTTGAAA 360
 30 CATAGGAGCT GCCCTCCTGG ATTTGGAGTG GTGCAAGCTG GAACCCCAAG GCGAAATACA 420
 GTTTGCAAAA GATGTCCAGA TGGGTTCTTC TCAAATGAGA CGTCATCTAA AGCACCCCTGT 480
 AGAAAACACA CAAATTGCAG TGTCTTTGGT CTCCTGCTAA CTCAGAAAGG AAATGCAACA 540
 CACGACAACA TATGTTCCGG AAACAGTGAA TCAACTCAAA AATGTGGAAT AGATGTTACC 600
 CTGTGTGAGG AGGCATTCTT CAGGTTTGCT GTTCCTACAA AGTTTACGCC TAACTGGCTT 660
 35 AGTGTCTTGG TAGACAATTT GCCTGGCACC AAAGTAAACG CAGAGAGTGT AGAGAGGATA 720
 AAACGGCAAC ACAGCTCACA AGAACAGACT TTCCAGCTGC TGAAGTTATG GAAACATCAA 780
 AACAAAGACC AAGATATAGT CAAGAAGATC ATCCAAGATA TTGACCTCTG TGAAAACAGC 840
 GTGCAGCGGC ACATTGGACA TGCTAACCTC ACCTTCGAGC AGCTTCGTAG CTTGATGGAA 900
 AGCTTACCGG GAAAGAAAGT GGGAGCAGAA GACATTGAAA AAACAATAAA GGCATGCAAA 960
 CCCAGTGACC AGATCCTGAA GCTGCTCAGT TTGTGGCGAA TAAAAAATGG CGACCAAGAC 1020
 40 ACCTTGAAGG GCCTAATGCA CGCACTAAAG CACTCAAAGA CGTACCACTT TCCCAAAACT 1080
 GTCACCTAGA GTCTAAAGAA GACCATCAGG TTCCTTCACA GCTTCACAA GTACAAATTG 1140
 TATCAGAAGT TATTTTGTAGA AATGATAGGT AACCAGGTCC AATCAGTAAA AATAAGCTGC 1200
 TTATAA 1206

Claims

1. A DNA comprising the nucleotide sequences of the Sequences No. 1 and No. 2 in the Sequence Table.
2. The DNA according to claim 1, wherein the Sequence ID No. 1 includes the first exon of the OCIF gene and the Sequence ID No. 2 includes the second, third, fourth, and fifth exons.
3. A protein exhibiting the activity of inhibiting differentiation and/or maturation of osteoclasts and having the following physicochemical characteristics,

(a) molecular weight (SDS-PAGE):

- (i) Under reducing conditions: about 60 kD,
- (ii) Under non-reducing conditions: about 60 kD and about 120 kD;

(b) amino acid sequence:

includes an amino acid sequence of the Sequence ID No. 3 in the Sequence Table,

(c) affinity:

exhibits affinity to a cation exchanger and heparin, and

(d) heat stability:

- (i) the osteoclastogenesis-inhibitory activity is reduced when treated with heat at 70°C for 10 minutes or at 56°C for 30 minutes,

- (ii) the osteoclastogenesis-inhibitory activity is lost when treated with heat at 90°C for 10 minutes.

4. A process for producing a protein exhibiting an activity of inhibiting differentiation and/or maturation of osteoclasts and having the following physicochemical characteristics,

(a) molecular weight (SDS-PAGE):

- (i) Under reducing conditions: about 60 kD,

- (ii) Under non-reducing conditions: about 60 kD and about 120 kD;

(b) amino acid sequence:

includes an amino acid sequence of the Sequence ID No. 3 of the Sequence Table,

(c) affinity:

exhibits affinity to a cation exchanger and heparin, and


(d) heat stability:

- (i) the osteoclastogenesis-inhibitory activity is reduced when treated with heat at 70°C for 10 minutes or at 56°C for 30 minutes,

- (ii) the osteoclastogenesis-inhibitory activity is lost when treated with heat at 90°C for 10 minutes,

the process comprising inserting a DNA including the nucleotide sequences of the sequences No. 1 and No. 2 in the Sequence Table into an expression vector, producing a vector capable of expressing a protein having the above-mentioned physicochemical characteristics and exhibiting the activity of inhibiting differentiation and/or maturation of osteoclasts, and producing this protein by a genetic engineering technique.

Figure 1



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP97/02859

A. CLASSIFICATION OF SUBJECT MATTER

Int. Cl⁶ C12N15/00, C12P21/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int. Cl⁶ C12N15/00, C12P21/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, GENETYX-CDROM, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Cancer Research, (1995), Vol. 55, Toshiyuki Yoneda, et al. "Sumarin suppresses hypercalcemia and osteoclastic bone resorption in nude mice bearing a human squamous cancer" P. 1989-1993	1 - 4
A	Proc. Natl. Acad. Sci. USA, (1990) Vol. 87 Kukita A. et al. "Osteoinductive factor inhibits formation of human osteoclast-like cells" P. 3023-3026	1 - 4

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

September 29, 1997 (29. 09. 97)

Date of mailing of the international search report

October 7, 1997 (07. 10. 97)

Name and mailing address of the ISA/

Japanese Patent Office

Facsimile No.

Authorized officer

Telephone No.